

References: Alberta Hip Fracture Clinical Pathway Acute Care

Last updated: February 1, 2012

Clinical Guidelines across the Care Continuum	1
Perioperative Care	2
Pain Management	3
Anticoagulation	3
Nausea	4
Infection	4
Skin and Wound Management	4
Osteoporosis	4
Delirium, Dementia and Depression	5
Nutrition	6
Rehabilitation	6
Falls	7
Care Transitions	8
Outcome Measures	8

Clinical Guidelines across the Care Continuum

Alberta Bone & Joint Health Institute. (July 2010). Evidence Review: Management of Hip Fracture. Calgary.

Beaupre, L.A., Jones, C.A., Majumdar, S.R., Johnston, D.W.C., Buckingham, J., & Saunders, L.D. State of the Science Review (2002): Continuum of Care and Delivery of Services for Elderly Patients with Hip Fracture. www.ahfmr.ab.ca/grants/state_of_science_final_reports.php 2002. Available from: URL: www.ahfmr.ab.ca/grants/state_of_science_final_reports.php

Beaupre, L.A., Jones, C.A., Saunders, L.D., Johnston, D.W.C., Buckingham, J., & Majumdar, S.R. Best Practices for Elderly Hip Fracture Patients: A Systematic Overview of the Evidence. *Journal of General Internal Medicine*, 2005; 20, 1019-25.

British Orthopaedic Association, British Geriatrics Society. Blue Book on Fragility Fractures. <http://www.fractures.com/pdf/BOA-BGS-Blue-Book.pdf> 7 A.D. September [cited 2011 Jun 10]; Available from: URL: <http://www.fractures.com/pdf/BOA-BGS-Blue-Book.pdf>

British Orthopaedic Association. (September, 2007). Standards for TRAUMA (BOAST). British Geriatrics Society.

Chilov, M., Cameron, I., & March, L. (2003). Evidence Based Guidelines for Fixing Broken Hips: an Update -Systematic Review. *Medical Journal of Australia*, 179, 489-493.

Mak, J. C., Cameron, I. D., & March, L. M. (2010). Evidence-based guidelines for the management of hip fractures in older persons: an update. *Medical Journal of Australia*, 192 (1), 37-41.

March, L., Chamberlain, A., Cameron, I., Cumming, R., Brnabic, A., Finnegan, T., et al. (1999, May 17). How best to fix a broken hip. *Medical Journal of Australia*, 170, pp. 489-494.

National Clinical Guideline Centre. (2011). The Management of Hip Fracture in Adults. Retrieved from www.ncgc.ac.uk

Neuman, M., Archan, S., Karlawish, J., Schwartz, J., & Fleisher, L. (November, 2009). The Relationship between Short-Term Mortality and Quality of Care for Hip Fracture: A Meta Analysis of Clinical Pathways for Hip Fracture. *The Geriatrics Society*, 2046-2054.

Oliver, D., Griffiths, R., Roche, J., & Sahota, O. (2007). Hip Fracture – Musculoskeletal Disorders. *BMJ Clinical Evidence*, 10 (1110), 1-27.

Scottish Intercollegiate Guidelines Network (SIGN). Management of hip fracture in older people. A national clinical guideline. <http://www.sign.ac.uk/pdf/sign111.pdf> 2009 June [cited 2011 Jun 6]; Available from: URL: <http://www.sign.ac.uk/pdf/sign111.pdf>

Waddell, J., McGlasson, R., Zellermeier, V., MacDonald, V., Lo, D. N., Spafford, D., et al. (June 30, 2011). National Hip Fracture Toolkit. The Bone and Joint Decade Canada. Alberta Bone & Joint Health Institute. (July 2010). Evidence Review: Management of Hip Fracture. Calgary.

Perioperative Care

Bjorkelund, K., Hommel, A., Thorngren, K., Lundberg, D., & Larsson, S. (February, 2011). The Influence of Perioperative Care and Treatment on the 4-Month Outcome in Early Patients with Hip Fracture. *AANA Journal*, 51-61.

Carson, J., Terrin, M., Noveck, H., Sanders, D., Chaitman, B., Rhoads, G., et al. Liberal or Restrictive Transfusion in High-Risk Patients after Hip Surgery. *The New England Journal of Medicine*, 365(26), 2453-62.

Leung, F., Lau, T., Khan, K., Chow, S., & Chow, A. (2010). Does Timing of Surgery Matter in Fragility Hip Fractures. *Osteoporosis International*, S529-S534.

Marsland, D., Colvin, P. L., Mears, S. C., & Kates, S. L. (September, 2010). How to optimize patients for geriatric fracture surgery. Baltimore; Rochester: International Osteoporosis Foundation and National Osteoporosis Foundation.

Waddell, J., McMullen, J., Lo, N., O'Connor, M., Sheppard, L., Mensour, M., et al. Improving Time to Surgery-Emergency Room, Preoperative and Immediate Postoperative Clinical Practice Guidelines for the Management of Hip Fracture Patients. <http://www.boneandjointhealthnetwork.ca> 2010 [cited 2011 Jun 6]; Available from: URL: <http://www.boneandjointhealthnetwork.ca>

Pain Management

Abou-Setta, A. M., Beupre, L. A., Rashid, S., Dryden, D. M., Hamm, M. P., Sadowski, C. A., et al. (May, 2011). Pain Management Interventions for Hip Fracture. Comparative Effectiveness Review No. 30. Prepared by the University of Alberta Evidence-based Practice Center under Contract No. 290-02-0023.) AHRQ Publication No. 11-EHC022-EF, available at: www.effectivehealthcare.ahrq.gov/reports/final.cfm, Edmonton.

Anticoagulation

Alberta Bone & Joint Health Institute. (Aug 2009). Anticoagulants Guidelines - Guidelines Review. Calgary.

Alberta Bone & Joint Health Institute. (Oct 2009). New Oral Anticoagulants - Evidence Review. Calgary.

American Academy of Orthopaedic Surgeons. (May, 2007). Prevention of Pulmonary Embolism in Patients Undergoing Total Hip or Knee Arthroplasty. Illinois: American Academy of Orthopaedic Surgeons Board of Directors.

Autar, R. (2010). A review of venous thromboprophylaxis in patients undergoing hip fracture surgery (HFS). *International Journal of Orthopaedic and Trauma Nursing*, 2-8.

Bell, A. D., Roussin, A., Cartier, R., Chan, W. S., Douketis, J. D., Gupta, A., et al. (2011). Society Guidelines: The use of Antiplatelet Therapy in Outpatient Setting: Canadian Cardiovascular Society Guidelines Executive Summary. *Canadian Journal of Cardiology*, 27 (Issue 2, March-April 2011), 208-221.

Chassot, P., Delabays, A., & Spahn, D. (2007, July 23). Perioperative antiplatelet therapy: the case for continuing therapy in patients at risk of myocardial infarction. *British Journal of Anaesthesia*, 99 (3), pp. 316-328.

Douketis, J. D. (2010, August 25). Pharmacologic Properties of the New Oral Anticoagulants: A Clinician-oriented Review with a Focus on Perioperative Management. *Current Pharmaceutical Design*, 16 (31), pp. 3436-3441.

Geerts, W. H., Bergqvist, D., Pineo, G. F., Heit, J. A., Samama, C. M., Lassen, M. R., et al. (2008). Prevention of venous thromboembolism: American College of Chest Physicians evidence-based clinical practice guidelines (8th Edition). *CHEST*, 33 (6 Suppl), 381S-453S.

Marsland, D., Mears, S., & Kates, S. (2010). Venous thromboembolic prophylaxis for hip fractures. *Osteoporosis International*, S593-S604.

NICE - National Institute for Health and Clinical Excellence. (January, 2010). Venous Thromboembolism: Reducing the Risk. London: NICE - National Institute for Health and Clinical Excellence.

Perka, C. (2011). Perioperative vs. postoperative initiation of thromboprophylaxis following major orthopaedic surgery: safety and efficacy of postoperative administration supported by recent trials of new oral anticoagulants. *Thrombosis Journal*, 9 (17), 1-20.

Ryn, J. v., Stangier, J., Haertter, S., Liesenfeld, K.H., Wienen, W., & Feuring, M. (2010). Dabigatran etexilate-a novel, reversible, oral direct thrombin inhibitor. *Thrombosis and Haemostasis*, 103 (6), pp. 1116-1127.

Safer Healthcare Now! (2010, Sept). Venous Thromboembolism Prevention in Hip Fracture Surgery. Retrieved from <http://www.saferhealthcarenow.ca/en/interventions/vte>

Nausea

Alberta Bone & Joint Health Institute. (Oct 2010). Evidence Review: Comparative Efficacy of Antiemetics. Calgary.

Carlisle, J. & Stevenson, C.A. (2009). Drugs for preventing postoperative nausea and vomiting. Cochrane Database of Systematic Reviews, vol. 1.

Gan, T. J., Meyer, T., Apfel, C. C., Chung, F., Davis, P. J., Eubanks, S., et al. (2003). Consensus Guidelines for Managing Postoperative Nausea and Vomiting. *Anesth Analg*, 97, pp. 62-71.

Perkins, P., Dorman, S. The Cochrane Collaboration. (2010). Haloperidol for the treatment of nausea and vomiting in palliative care patients (Review). *The Cochrane Collaboration*, 1-12.

Infection

Alberta Bone & Joint Health Institute. (Aug 2009). Evidence Review: Prophylactics Antibiotic Use in Hip and Knee Arthroplasty.

Safer Healthcare Now! (Mar, 2011). *Prevent Surgical Site Infections: Getting Started Kit*. Retrieved from <http://www.saferhealthcarenow.ca/en/interventions/ssi>

Skin and Wound Management

Braden, B., Bergstrom, N. Braden Scale for Predicting Pressure Sore Risk. [Cited 2011 Jun 10]; Available from: URL: <http://www.bradenscale.com/images/bradenscale.pdf>

Canadian Agency for Drugs and Technologies in Health (CADTH). (2011). Rapid Response Report: Surgical Dressing for Patients Undergoing Hip and Knee Arthroplasty: Clinical Effectiveness and Cost Effectiveness. CADTH.

Kring, D.L. Reliability and validity of the Braden Scale for predicting pressure ulcer risk. *Journal of Wound, Ostomy & Continence Nursing* 2007 Jul; 34(4):399-406.

Osteoporosis

Alberta Bone & Joint Health Institute. (Aug 2009). Evidence Report: Osteoporosis Management in Hip Fracture Patients. Calgary.

Beaupre, L.A., Morrish, D.W., Hanley, D.A., Maksymowych, W.P., Bell, N.R., Juby, A.G., Majumdar, S.R. Oral bisphosphonates are associated with reduced mortality after hip fracture. *Osteoporosis International*. 2011 Mar; 22(3):983-91. Epub 2010 Nov 4.

Bischoff-Ferrari, H.A., Dawson-Hughes, B., Platz, A., Orav, E.J., Stahelin, H.B., Willett, W.C., et al. Effect of high-dosage cholecalciferol and extended physiotherapy on complications after hip fracture: a randomized controlled trial. *Arch.Intern.Med.* 2010 May 10; 170(9):813-820.

Bischoff-Ferrari, H.A., Willett, W.C., Wong, J.B., Stuck, A.E., Staehelin, H.B., Orav, E.J., et al. Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. *Arch.Intern.Med.* 2009 Mar 23; 169(6):551-561.

Blouin, J., et al. Discontinuation of Antiresorptive Therapies: A Comparison between 1998–2001 and 2002–2004 among Osteoporotic Women. *J Clin Endocrinol Metab* 2007; 92:887-94. (Canadian Study)

Grey, A., Bolland, M., Wattie, D., Horne, A., Gamble, G., Reid, I.R. Prolonged antiresorptive activity of zoledronate: a randomized, controlled trial. *J Bone Miner Res*. 2010 Oct; 25(10):2251-5.

Grima, D.T., Papaioannou, A., Airia, P., Ioannidis, G., Adachi, J.D. Adverse events, bone mineral density and discontinuation associated with generic alendronate among postmenopausal women previously tolerant of brand alendronate: a retrospective cohort study. *BMC Musculoskelet Disord*. 2010 Apr 14; 11:68. (Canadian Study)

Hanley, D.A, Cranney, A., Jones, G., Whiting, S.J., Leslie, W.D., Cole, D.E.C., et al. Vitamin D in adult health and disease: a review and guideline statement from Osteoporosis Canada. *CMAJ* 2010 Sep 7; 182(12):E610-E618.

Lyles, K.W., Colon-Emeric, C.S., Magaziner, J.S., et al. HORIZON Recurrent Fracture Trial. Zoledronic acid and clinical fractures and mortality after hip fracture. *N Engl J Med* 2007; 357: 1799-1809.

Majumdar, S., Beaupre, L., Harley, C., Hanley, D., Lier, D., Juby, A., et al. (2007). Use of a Case Manager to Improve Osteoporosis Treatment after Hip Fracture. *Archives of Internal Medicine*, 167(19), 2110-2115.

Miller, P.D., et al. Safety and efficacy of risedronate in patients with age-related reduced renal function as estimated by the Cockcroft and Gault method: a pooled analysis of nine clinical trials. *J Bone Miner Res*. 2005 Dec; 20(12):2105-15.

Papaioannou, A., Morin, S., Cheung, A.M., Atkinson, S., Brown, J.P., Feldman, S., et al. 2010 clinical practice guidelines for the diagnosis and management of osteoporosis in Canada: summary. *Canadian Medical Association Journal* 2010 Nov 23; 182(17):1864-73.

Rahmani, P., & Morin, S. (November, 2009). Prevention of Osteoporosis-Related Fractures among Postmenopausal Women and Older Men. *CMAJ - Review*, 815-820.

Ringe, J.D., Möller, G. Differences in persistence, safety and efficacy of generic and original branded once weekly bisphosphonates in patients with postmenopausal osteoporosis: 1-year results of a retrospective patient chart review analysis. *Rheumatol Int*. 2009 Dec; 30(2):213-21.

Siris, E.S., Harris, S.T., Rosen, C.J., et al. Adherence to bisphosphonate therapy and fracture rates in osteoporotic women: relationship to vertebral and nonvertebral fractures from 2 US claims databases. *Mayo Clin Proc* 2006; 81:1013-22.

Tosi, L.L., Gliklich, R., Kannan, K., Koval, K.J. The American Orthopaedic Association's "own the bone" initiative to prevent secondary fractures. *J Bone Joint Surg.Am*. 2008 Jan; 90(1):163-173.

Delirium, Dementia and Depression

Holroyd-Leduc, J. M., Khandwala, F., & Sink, K. M. (2009, August 17). How can delirium best be prevented and managed in older patients in hospitals? *CMAJ - Canadian Medical Association or its Licensors*, pp. 1-6.

Inouye, S.K. Delirium After Hip Fracture: To Be or Not to Be? *Journal of the American Geriatrics Society* 2001; 49(5):678-679.

McGilton, K., Wells, J., Davis, A., Rochon, E., Calabrese, S., Teare, G., et al. Rehabilitating patients with dementia who have had a hip fracture: Part II: cognitive symptoms that influence care. *Topics in Geriatric Rehabilitation* 2007; 23(2):174-82.

McGilton, K., Wells, J., Teare, G., Davis, A., Rochon, E., Calabrese, S., et al. Rehabilitating patients with dementia who have had a hip fracture - Part I: Behavioral symptoms that influence care. *Topics in Geriatric Rehabilitation* 2007; 23(2):161-73.

McGilton, K.S., Mahomed, N., Davis, A.M., Flannery, J., Calabrese, S. Outcomes for older adults in an inpatient rehabilitation facility following hip fracture (HF) surgery. *Archives of Gerontology & Geriatrics* 2009 Jul; 49(1):e23-e31.

NHS National Institute for Health and Clinical Excellence. (2010, July). Delirium: Diagnosis, Prevention and Management. NICE Clinical Guideline 103, pp. 1-29.

Nutrition

Avenell, A., Handoll, H.C.G. (2010). Nutritional Supplementation for Hip Fracture Aftercare in Older People (Review). *Cochrane Database Syst Rev*, 1-86.

Dillabough, A., Mammel, J., & Yee, J. (2011). Improving nutritional intake in post-operative hip fracture patients: A quality improvement project. *International Journal of Orthopaedic and Trauma Nursing*, 196-201.

Fabian, E., Gerstorfer, I., Thaler, H., Stundner, H., Biswas, P., & Elmadfa, I. (2011, January 21). Nutritional Supplementation Affects Postoperative Oxidative Stress and Duration of Hospitalization in Patients with Hip Fracture. *The Central European Journal of Medicine*, pp. 88- 93.

Hommel, A., Bjorkelund, K.B., Thorngren, K.G., Ulander, K. Nutritional status among patients with hip fracture in relation to pressure ulcers. *Clinical Nutrition* 2007 Oct; 26(5):589-96.

Rizzoli, R. (2011). Nutritional Needs and Interventions to Support Geriatric Hip Fracture Recovery. *Clinical Nutritional Highlights*, 7 (2), 2-8.

Schurch, M.A., Rizzoli, R., Slosman, D., Vadas, L., Vergnaud, P., & Bonjour JP. (1998). Protein Supplements Increase Serum Insulin-Like Growth Factor-I Levels and Attenuate Proximal Femur Bone Loss in Patients with Recent Hip Fracture: A Randomized, Double-Blind, Placebo-Controlled Trial. *Ann Intern Med*, May 15, 1998 128:801-809.

Rehabilitation

Bachmann, S. Inpatient rehabilitation specifically designed for geriatric patients: systematic review and meta-analysis of randomised controlled trials. *British Medical Journal* 2010; 340.

Beaupre, L.A., Cinats, J.G., Jones, C.A., Scharfenberger, A., Johnston, D.W.C., Senthilselvan, A., et al. Does Functional Recovery in Elderly Hip Fracture Patients differ between Patients Returning to Long-Term Care or the Community? *The Journals of Gerontology. Series A. Biological sciences and Medical Sciences*.2007; 62(10):1127-33.

Beaupre, L.A., Cinats, J.G., Senthilselvan, A., Lier D, Jones CA, Scharfenberger A, et al. Reduced Morbidity for Elderly Patients with Hip Fracture after Implementation of a Perioperative Evidence- Based Clinical Pathway. *Quality and Safety in Health Care* 2006; 15(5): 375–379.

Beaupre, L.A., Cinats, J.G., Senthilselvan, A., Scharfenberger, A., Johnston, D.W.C., Saunders, L.D. Does Standardized Rehabilitation and Discharge Planning Improve Functional Recovery in Elderly Patients with Hip Fracture? *Archives of Physical Medicine & Rehabilitation* 2005; 86(12):2231-9.

Beaupre, L.A., Jones, C.A., Saunders, L.D., Johnston, D.W.C., Buckingham, J., Majumdar, S.R. Best Practices for Elderly Hip Fracture Patients: A Systematic Overview of the Evidence. *Journal of General Internal Medicine* 2005; 20:1019-25.

Bellelli, G., Frisoni, G.B., Pagani, M., Magnifico, F., Trabucchi, M. Does cognitive performance affect physical therapy regimen after hip fracture surgery? *Aging Clin Exp Res* 2007; 19(2):119-24.

Chudyk, A.M., Jutai, J.W., Petrella, R.J., Speechley, M. Systematic review of hip fracture rehabilitation practices in the elderly. *Archives of Physical Medicine & Rehabilitation* 2009; 90(2):246-62.

Halbert, J., Crotty, M., Whitehead, C., Cameron, I., Kurrle, S., Graham, S., et al. Multi-disciplinary rehabilitation after hip fracture is associated with improved outcome: A systematic review. *Journal of Rehabilitation Medicine* 2007; 39(7):507-12.

Handoll, H.H. Multidisciplinary rehabilitation for older people with hip fractures. 2009 Cochrane Database of Systematic Reviews (4):CD007125.

Handoll, H., Sherrington, C., & Mak, J. (2011). Interventions for Improving Mobility after Hip Fracture Surgery in Adults - Intervention Review. The Cochrane Library.

Healee, D., McCallin, A., & Jones, M. (2011). Older Adult's Recovery from Hip Fracture: A Literature Review. *International Journal of Orthopaedic and Trauma Nursing*, 18-28.

McGilton, K.S. Outcomes for older adults in an inpatient rehabilitation facility following hip fracture (HF) surgery. *Archives of Gerontology and Geriatrics* 2009 Jul; 49(1):e23-e31.

Mehta, S., & Roy, J. (2011). Systematic Review of Home Physiotherapy after Hip Fracture Surgery. *J Rehabil Med*, 477-480.

Falls

Cochrane Review: Interventions for Preventing Falls in Older People Living in the Community: 2009.

Cochrane Review: Interventions for Preventing Falls in Older People in Nursing Care Facilities and Hospitals: 2010.

Di Marco, M., Vallero, F., deToma, E., deLauso, L., Tappero, R, Cavanna, A. A Single Home Visit by an Occupational Therapist Reduces the Risk of Falling after Hip Fracture in Elderly Women: a quasi-randomized controlled trial. *Journal of Rehabilitation Medicine*. 2008; 40:446-450.

Gates, S., Lamb, S. E., Fisher, J. D., Cooke, M. W., & Carter, Y. H. (2008). Multifactorial Assessment and Targeted Intervention for Preventing Falls and Injuries among Older People in Community and Emergency Care Settings: Systematic Review and Meta-Analysis. Warwick: *BMJ*.

Gillespie, L., Robertson, M., Gillespie, W., Lamb, S., Gates, S., Cumming, R., et al. (2009, Apr 15). Interventions for Preventing Falls in Older People Living in the Community (Review). The Cochrane Collaboration, 1-322.

Holroyd-Leduc, J. (December, 2009). Therapeutics - Review: Exercise Interventions Reduce Falls in Elderly People Living in the Community. *Evidence Based Medical Journal*, 176.

Rahme, E., Kahm, S.R., Dasgupta, K. et al. Short-term mortality associated with failure to receive home care after hemiarthroplasty. *Canadian Medical Association Journal*. 2010; 182:1421-1426.

Rose, D. Preventing Falls Among Older Adults No "one size suits all" Intervention Strategy. *JRRD*. 2008; 8:1153-1166.

Russell, M.A., Hill, K.D., Blackberry, I., Day, L.L., Dharmage, C.S. Falls risk and functional decline in older fallers discharged directly from emergency departments. *Journal of Gerontology*. 2006; 61A:1090-95.

Sherrington, C., Lord, S.R., Close, J.C.T. Best Practice Recommendations for Physical Activity to Prevent Falls in Older Adults: 2008. Accessed December 3, 2010. <http://fallsnetwork.powmri.edu.au/resources/files/aaa-sax-pa-falls.pdf>

Teppo et al, Shifting the Focus in Fracture Prevention from Osteoporosis to Falls: *British Medical Journal*, 2008; 336:124-126.

Visschedijk, J., Achterberg, W., van Balen, R., Hertogh, C. Fear of Falling after Hip Fracture: A systematic review of measurement instruments, prevalence, interventions, and related factors. *Journal of the American Geriatric Society*. 2010; 58:1739-48.

Care Transitions

Blaylock, A., & Cason, C. (July, 1992). Discharge Planning - Predicting Patients Needs. *Journal of Gerontological Nursing*, 5-10.

Rahme, E., Kahn, S., Dasgupta, K., Berman, M., Bernatsky, S., Habel, Y., et al. (September, 2010). Short-Term Mortality Associated with Failure to Receive Home Care after Hemiarthroplasty. *CMAJ Research*, 1421-1426.

Outcome Measures

Bryant, D., Sanders, D., Coles, C., Petrisor, B., Jeray, K., & Laflamme, F. (2009, November 6). Selection of Outcome Measures for Patients with Hip Fracture. *J Orthop Trauma*, pp. 434-441.