

CONTACT PRECAUTIONS SPORICIDAL CLEAN



CLEAN HANDS
UPON ENTRY

SOAP AND WATER
RECOMMENDED
FOR HAND HYGIENE
AFTER REMOVING
GLOVES



STAFF AND VISITORS



VISITORS:
GOWN AND GLOVE ONLY
IF ASSISTING WITH CARE.

PLEASE CHECK WITH
NURSING STAFF
BEFORE ENTERING
ROOM.

PATIENTS UPON LEAVING ROOM



(FOR ESSENTIAL PURPOSES ONLY)
(CLEAN GOWN/CLOTHES AND HANDS)

**SINGLE ROOM RECOMMENDED
WITH DEDICATED EQUIPMENT**

CONTACT PRECAUTIONS

SPORICIDAL CLEAN

Acute and Continuing Care

Use in Addition to Routine Practices

Additional Information

- **Soap and water recommended for hand hygiene when providing care for patients with diarrhea.** (AHS Hand Hygiene Procedure, June 2017)
- Communicate with Environmental Services the need for sporicidal cleaning procedures:
 - All high touch surfaces in the bedroom and bathroom must receive a daily sporicidal clean (**ES-PROT-CLN-061**)
 - Sporicidal Discharge/Transfer Clean protocol must be followed when a patient is
 - discharged,
 - transferred to a new room or unit,
 - transferred to another facility, or
 - remaining in the room AND sporicidal precautions are discontinued.
(**ES-PROT-CLN-060**)
 - Sign is to remain posted until Discharge/Transfer Sporidal clean is completed.
- Dedicate equipment to the isolated patient or clean and disinfect shared equipment after every use (to prevent transmission).
- Patient to leave room only for **essential** purposes. If patients are leaving their room they should be cognitively intact or supervised, have clean hands and clothing and any drainage and/or body fluids contained.

Common Organisms Requiring SPORICIDAL Precautions (not all inclusive)

- *Clostridium difficile* (CDI)
- *Candida auris* multidrug-resistant