

Triage Letter - take this letter with you to the Emergency Department and give it to the Nurse or Doctor when you arrive.

To: Emergency Department Physician/Nurse

RE: CHECKPOINT INHIBITOR ADVERSE REACTION

From: Dr	(patient's oncologist) at
(cancer cer	ntre).
This patient has received treatm	ent with a Checkpoint Inhibitor called:
	_

Immune-related side effects are known to occur following this treatment. Side effects occur due to unintended tissue injury following treatment with immunotherapy, and any organ system may be involved.

These immune-related side-effects may be severe or life-threatening, and require different management from chemotherapy associated toxicities.

If immune-related side-effects are suspected, **contact the medical oncologist listed above (or the covering designate).** With prompt management, immune-related side-effects are generally reversible, however immunosuppression may be required.

Signs and symptoms suggestive of immune-related side-effects:				
Gastrointestinal: ◆ Colitis	Diarrhea Nausea Elevated lipase	Gastrointestinal bleedingElevated liver enzymesAbdominal pain/cramping		
Pulmonary: ◆ Pneumonitis	Shortness of breathCough	Chest painFever		
Skin: ◆ Dermatitis	Pruritic rash			
Endocrine:	Fatigue, lethargyDehydrationHypotension	Headache, visual disturbancesMental status changesAnxiety, nervousness, tremors		
Musculoskeletal: ◆ Arthritis ◆ Myositis	Joint pain/swelling Fatigue/myalgias			
Neurological: ◆ Peripheral neuropathy ◆ CNS inflammation	Numbness/pain Mental status change	Headache Focal neurological deficit		

Signs and symptoms suggestive of immune-related side-effects:				
Renal: ◆ Acute kidney injury	Swelling Elevated serum creatinine			
Cardiac: ◆ Myocarditis ◆ Pericarditis	Chest pain Fever	ECG changesElevated cardiac enzymes		
Fever: 38.9 °C or higher with any of the following:	 lasts longer than 24 hours feels unwell	 with other symptoms: cough, shortness of breath, pain, dizziness, nausea or diarrhea 		
*If the patient is also taking chemotherapy, follow the chemotherapy fever guidelines instead:	They have been told to report to the nearest ER as soon as possible if febrile (a single oral temperature of 38.3°C/100.9°F or higher or an oral temperature reading of 38.0°C/100.4°F measured at 2 occasions, 1 hour apart). If left untreated, this complication of febrile neutropenia could lead to life-threatening sepsis.			

For complete recommendations and evidence, access the full guideline at:

www.ahs.ca/guru > Symptom Management > Checkpoint Inhibitor Toxicities

CONTACT INFORMATION:

Daytime Weekdays		
Calgary (Arthur Child)	Call patient's oncologist/hematologist as listed above directly or call 587-231-3131 and ask to speak to the patient's oncologist/hematologist.	
	After hours, weekends and holidays call 587-231-3100, press 0, and request the on-call oncologist/hematologist, or use ROCA	
Edmonton (Cross Cancer Institute)	780-432-8771, After hours, weekends and holidays - contact the on-call oncologist/hematologist, or use ROCA.	
Grande Prairie (Grande Prairie Cancer Centre)	825-412-4200	
Lethbridge (Jack Ady Cancer Centre)	403-388-6802	
Medicine Hat (Margery E. Yuill Cancer Centre)	403-529-8817	
Red Deer (Central Alberta Cancer Centre)	403-343-4526	
After Hours		
RAAPID North at 1-800-282-9911		
RAAPID South at 1-800-661-1700		

Thank you for your assistance in the care of this patient