How to Prepare for Your Prostate Brachytherapy Implant

Cross Cancer Institute

This is a companion resource to the "Prostate Brachytherapy" booklet and will give you specific information on how to prepare for this procedure.



Once you know your brachytherapy appointment date:



Please follow the instructions below carefully. Start 2 weeks (14 days) before your implant.



Arrange for someone to drive you to your appointment and pick you up. It is not safe to drive after you have anesthetic. The Pre-Admission Clinic will give you more information.

2 Weeks Before Implant	 Buy from your local pharmacy: Fleet Enema[®], no prescription needed Ciprofloxacin[®] antibiotic (with your prescription) Flomax[®], also known as Tamsulosin (with your prescription) Stop taking vitamins or herbal supplements (you can still take Vitamin D and Calcium)
1 Week Before Implant	 Stop taking blood thinners: If you are taking Aspirin[®], Ibuprofen[®], or drugs that contain Aspirin[®], you will start taking these medications again 1 week after the implant. If you are on any other prescription blood thinners, you will have an appointment with an anesthesiologist who will give you instructions on how to manage this medication. Start taking Flomax[®]. Take 1 pill in the evening
Weekend Before Implant	☐ Start low fibre diet as directed by your brachytherapist
1 Day Before Implant	 Eat and drink throughout the day. Make sure you get enough fluids so you are fully hydrated for your surgery. You can drink clear fluids such as water, apple juice, cranberry juice or black coffee Avoid dairy or alcohol Start taking Ciprofloxacin® as prescribed - 1 pill in the morning and 1 pill in the evening. Phone 780-432-8951 between 1 – 3pm to confirm your arrival time at the Cross Cancer Institute. If the day before your surgery is a holiday, then call the last business day before your surgery. Stop eating after midnight (12 am). You may drink water until 5:00 am.

Radiation Treatment

	Starting at midnight (12 am) before the procedure:
Day of Implant Before procedure	Starting at miding it (12 and before the procedure.
	□ From midnight to 5:00 am, drink only water
	 On not drink anything after 5:00 am You may take small sips of water if you need to take medication
	 Not may take small sips of water if you need to take medication O not take a Ciprofloxacin[®] pill on the morning of the implant. You will get your morning dose of Ciprofloxacin[®] in the operating room by IV (intravenous) while you are asleep Give yourself a Fleet Enema[®] at least 1 hour before you leave for the
	Cross Cancer Institute
	Going to the hospital:
	\Box Bring your healthcare card and government ID
	🛞 Do not bring any valuables
	Bring along activities to help you pass the time and any medication or other things you need for the day. You will be at the hospital for several hours
	At the hospital:
	Check in at the Main Reception Desk
	Your healthcare team will prepare you for your implant
	You will meet the Anesthesiologist, Radiation Oncologist and O.R. Nurse once you are in the operating room at the Cross Cancer Institute
The procedure	The procedure will take about 1 hour and 30 minutes
Day of Implant After procedure	After the implant you will get a radiation safety Blue Wallet Card and discharge instructions
	☐ Take your Ciprofloxacin [®] pill in the evening. Continue to take as
	prescribed until your pills are done
	□ Take your Flomax [®] in the evening
Follow-up Appointment	You will have a follow-up appointment in person or by phone about 4 weeks after the implant with your Radiation Oncologist
Questions about Treatment?	
Brachytherapy Office 780-432-8475	
	Monday to Friday, 7:30 am – 4:00 pm
	www.cancercarealberta.ca

1)