Routine Immunization Schedule

A		
Age	Vaccine(s) your child/you will get	Diseases your child/you will be protected against
2 months	 DTaP-IPV-Hib-HB * Pneumococcal conjugate Rotavirus 	 Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b, hepatitis B Pneumococcal Disease Rotavirus
4 months	 DTaP-IPV-Hib-HB * Pneumococcal conjugate Meningococcal conjugate (MenconC) Rotavirus 	 Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b, hepatitis B Pneumococcal Disease Meningococcal Disease Rotavirus
6 months	 DTaP-IPV-Hib-HB * Pneumococcal conjugate (for children at high risk only) 	• Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b, hepatitis B
12 months	 MMR-Var Pneumococcal conjugate Meningococcal conjugate (MenconC) 	 Measles, mumps, rubella, varicella (chickenpox) Pneumococcal Disease Meningococcal Disease
18 months	DTaP-IPV-HibMMR-Var	 Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b Measles, mumps, rubella, varicella (chickenpox)
4 years	• Tdap-IPV	 Tetanus, diphtheria, pertussis (whooping cough), polio
Grade 6	 Hepatitis B (2 doses, 6 months apart if not already given in childhood) HPV (2 or 3 [†] doses over 6 months) 	Hepatitis BHuman papillomavirus
Grade 9	TdapMenC-ACYW	 Tetanus, diphtheria, pertussis (whooping cough) Meningococcal Disease

* DTaP-IPV-Hib-HB – for children born on or after March 1, 2018. Children born before March 1, 2018, are offered DTaP-IPV-Hib at ages 2, 4 and 6 months and Hepatitis B vaccine in school.
 † A 3rd dose of HPV is needed only for students starting the vaccine series at 15 years of age

or older and anyone with a weak immune system.

See reverse for more information.



Age	Vaccine(s) your child/you will get	Diseases your child/you will be protected against
Adult	 Tdap (1 dose for adults who are not immunized for pertussis and 1 booster dose every 10 years; if pregnant see below) Hepatitis B (3 doses for unprotected adults born in 1981 or later) HPV (3 doses for adults up to and including age 26 years who did not get HPV vaccine in school) MMR for adults born in 1970 or later and no record of vaccine (2 doses) Varicella (chickenpox) for unprotected adults (2 doses) Pneumococcal conjugate at 65 years of age or older Polio (3 doses for people who are not immunized) RSV (1 dose for adults age 60 years and older who live in continuing care and supportive living facilities or are Indigenous, and adults age 70 years and older) 	 Tetanus, diphtheria, pertussis (whooping cough) Hepatitis B Human papillomavirus Measles, mumps, rubella Varicella (chickenpox) Pneumococcal Disease Polio Respiratory syncytial virus
Pregnant	• Tdap with every pregnancy	• Tetanus, diphtheria, pertussis (whooping cough)
Everyone (6 months of age and older)	 Annual influenza (2 doses for children under age 9 years if getting this vaccine for the first time) COVID-19[‡] 	InfluenzaCOVID-19

When to get a COVID-19 vaccine depends on your age, health, and when you had your last dose. Go to <u>ahs.ca/covidvaccine</u> for more information.

Note:

More vaccines may be needed because of health conditions, work, school, living arrangements, household contacts, lifestyle risks, travel, or because a person was not fully immunized in childhood. Check with your healthcare provider, a public health nurse, your workplace health and safety department, or a travel health clinic to find out if you need any other vaccines and if they are free for you.

For more information about immunization



Call Health Link at 811



Scan the QR code or go to ahs.ca/immunize

Adult Immunization

