

Postpartum Depression Education Module: References by Topic Area

The following articles can be ordered from Alberta Health Services Knowledge Resource Services by requesting an article at <https://krs.libguides.com/articlerequest>

Key Guiding Documents

American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-5. Washington, DC; 2013.

Austin M, Hight N, Group EW. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia; 2017.

Austin M, Hight N, Guidelines Expert Advisory Committee. Clinical practice guidelines for depression and related disorders – anxiety, bipolar disorder and puerperal psychosis – in the perinatal period. A guideline for primary care health professionals. [Internet]. Melbourne, Australia; 2011.

BC Reproductive Mental Health Program & Perinatal Services. Best Practice Guidelines for Mental Health Disorders in the Perinatal Period. [Internet]. 2014.

National Collaborating Centre for Mental Health Royal College of Psychiatrists' Research and Training. Antenatal and Postnatal Mental Health: The NICE guideline on Clinical Management and Service Guidance. Updated edition. Leicester; 2014

(SIGN) Network SIG, Scottish Intercollegiate Guidelines Network (SIGN). Management of perinatal mood disorders [Internet]. Edinburgh; 2012.

Acceptability, Barriers, Stigma and Help seeking

Austin M. Antenatal screening and early intervention for perinatal distress depression and anxiety: where to from here? Arch Women's Ment Heal. 2004; 7:1–6.

Bowen A, Bowen R, Butt P, Rahman K, Muhajarine N. Patterns of depression and treatment in pregnant and postpartum women. Can J Psychiatry. 2012; 57(3):161–7.

Buist A, Bilszta J. The beyondblue National Postnatal Screening Program, Prevention and Early Intervention 2001-2005, Final Report. Vol 1: National Screening Program. Melbourne; 2006.

Dennis C, Chung-Lee L. Postpartum depression help-seeking barriers and maternal treatment preferences: a qualitative systematic review. Birth. 2006; 33(4):323–31.

Hight N, Stevenson A, Purtell C. Qualitative insights into women's personal experiences of perinatal depression and anxiety. Women Birth. 2014; 27(3):179–84.

Gemmill A, Leigh B, Eriksen J. A survey of clinical acceptability of screening for postnatal depression in depressed and non-depressed women. BMC Public Health. 2006; 6:211.

References by Topic Area | 2

- Kingston D, Austin M, Heaman M. Barriers and facilitators of mental health screening in pregnancy. *J Affect Disord*. 2015; 186:350–7.
- Kingston D, Biringer A, McDonald S. Preferences for mental health screening among pregnant women: a cross-sectional study. *Am J Prev Med*. 2015; 49(4):e35–43.
- Kingston D, McDonald S, Tough S, Austin M-P, Hegadoren K, Lasiuk G. Public views of acceptability of perinatal mental health screening and treatment preference: a population based survey. *BMC Pregnancy Childbirth*. 2014 Dec; 14(1):67.
- Liberto T. Screening for depression and help-seeking in postpartum women during well-baby pediatric visits: an integrated review. *J Pediatr Heal Care [Internet]*. 2012; 26(2):109–17.
- Megnin-Viggars O, Symington I, Howard L. Experience of care for mental health problems in the antenatal or postnatal period for women in the UK: a systematic review and meta-synthesis of qualitative research. *Arch Womens Ment Heal*. 2015; 18(6):745–59.
- Moore D, Ayers S, Drey N. A Thematic Analysis of Stigma and Disclosure for Perinatal Depression on an Online Forum. *JMIR Ment Heal*. 2016; 3(2).
- National Institute for Health Care Management. Identifying and treating maternal depression: strategies and considerations for health plans. [Internet]. Washington; 2010.
- Sharma V, Sharma P. Postpartum depression: diagnostic and treatment issues. *JOGC*. 2012; 34(5):436–42
- Teng L, Blackmore Robertson E, Stewart D. Healthcare worker’s perceptions of barriers to care by immigrant women with postpartum depression: an exploratory qualitative study. *Arch Women’s Ment Heal*. 2007; 10(3):93–101

Clinical Judgment

- Tanner C. Thinking like a nurse: a research-based model of clinical judgment in nursing. *J Nurs Educ [Internet]*. 2006; 45(6):204–2011.

Cultural Considerations for PPD for Indigenous Peoples

- Daoud N, Smylie J. Unpublished analysis, Canadian Maternity Experiences Survey. Toronto, Ontario; 2014.
- Indigenous Health Program. Indigenous Health Program Telehealth Learning Series [Internet]. 2015 [cited 2016 Nov 19].
- Indigenous Health Program. Practical Considerations Working with Aboriginal Mothers. Alberta Health Services; 2016.
- National Aboriginal Health Organization (NAHO). Celebrating birth-Aboriginal midwifery in Canada. Ottawa, Ontario; 2008.

Reading C, Wien F. Health inequalities and the social determinants of Aboriginal peoples' health [Internet]. Prince George, BC; 2009.

Smylie J. Strong Women, Strong Nations. Aboriginal Maternal Mental Health in British Columbia [Internet]. Toronto, Ontario; 2014.

Diversity and Cultural Considerations for PPD in Diverse Populations

Calgary Health Region. Communicating with Diverse Populations. 2008.

Cross T, Bazron B, Dennis K, Isaacs M. Towards a Culturally Competent System of Care, Volume 1. Washington, DC; 1989.

Hight N, Stevenson A, Purtell C. Qualitative insights into women's personal experiences of perinatal depression and anxiety. *Women Birth*. 2014; 27(3):179–84.

Kingston D, Austin M, Heaman M. Barriers and facilitators of mental health screening in pregnancy. *J Affect Disord*. 2015; 186:350–7.

Michigan Department of Community Health Behavioral Health and Developmental Disabilities Administration. Transforming Cultural and Linguistic Theory into Action: A Toolkit for Communities [Internet]. 2012.

Moore D, Ayers S, Drey N. A Thematic Analysis of Stigma and Disclosure for Perinatal Depression on an Online Forum. *JMIR Ment Heal*. 2016;3(2).

National Center for Cultural Competence Georgetown University Centre for Child and Human Development. Curricula Enhancement Module Series Definitions of Cultural Competence. [Internet]. 2012 [cited 2016 Mar 29].

Primary Health Care Programs Alberta Health Services. Diversity Awareness Self-Reflection Tool - A Practical Tool for Health Care Teams [Internet]. 2015.

Registered Nurses' Association of Ontario. Embracing Cultural Diversity in Health Care: Developing Cultural Competence [Internet]. Toronto, Canada; 2007.

Teng L, Blackmore Robertson E, Stewart D. Healthcare worker's perceptions of barriers to care by immigrant women with postpartum depression: an exploratory qualitative study. *Arch Women's Ment Heal*. 2007; 10(3):93–101.

Family Support Plan

Austin M, Hight N, Group EW. *Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline*. Melbourne, Australia; 2017.

BC Reproductive Mental Health Program & Perinatal Services. *Best Practice Guidelines for Mental Health Disorders in the Perinatal Period*. [Internet]. 2014.

National Collaborating Centre for Mental Health Royal College of Psychiatrists' Research and Training. *Antenatal and Postnatal Mental Health: The NICE guideline on Clinical Management and Service Guidance*. Updated edition. Leicester; 2014.

Impact of Maternal Mental Illness

Austin M, Hight N, Group EW. *Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline*. Melbourne, Australia; 2017.

Beck C. Postpartum Depression. It isn't just the blues. *AJN*. 2006; 106(5):40–50.

Beck C. Maternal depression and child behaviour problems: a meta-analysis. *J Adv Nurs*. 1999; 29(3):623–9.

Cox J, Holden J, Henshaw C. *Perinatal Mental Health: The Edinburgh Postnatal Depression Scale (EPDS) Manual*. 2nd ed. RC Psych Publications; 2014.

Dennis CL MK. The relationship between infant-feeding outcomes and postpartum depression: A qualitative systemic review. *Pediatrics*. 2009; 123(4):e736–51.

Goodman J. Paternal depression, its relationship to maternal depression and implications for family health. *J Adv Nurs*. 2004; 45(1):26–35.

Horwitz S, Briggs-Gowan M, Storfer-Isser A, Carter A. Persistence of maternal depressive symptoms throughout the early years of childhood. *J Womens Heal*. 2009; 18(5):637–45.

Kingston D, Tough S, Whitfield H. Prenatal and postpartum maternal psychological distress and infant development: a systematic review. *Child Psychiatry Hum Dev*. 2012; 43(5):683–714.

Moehler E, Brunner R, Wiebel A, Reck C, Resch F. Maternal depressive symptoms in the postnatal period are associated with long-term impairment of mother-child bonding. *Arch Womens Ment Heal*. 2006; 9(5):273–8.

Murray L, Cooper P. Postpartum depression and child development. *Psychol Med*. 1997; 27(2):253–60.

Perinatal Mental Health Consortium. *National Action Plan for Perinatal Mental Health 2008–2010 Full Report*. [Internet]. Melbourne, Australia; 2008.

Priest S, Barnett B. Perinatal anxiety and depression: issues, outcomes and interventions. In: *Infants of Parents with Mental Illness: Developmental, Clinical, Cultural and Personal Perspectives*. Bowen: Hills Australian Academic Press; 2008.

Mental Health Concepts

Canadian Mental Health Association Alberta. *Mental Illness in Canada*.

Canadian Mental Health Association British Columbia. *Stigma and Discrimination around mental health and substance use problems* [Internet]. British Columbia; 2014.

Canadian Mental Health Association. *Mental Health for Life* [Internet]. 2016.

Centre for Chronic Disease Prevention Public Agency of Canada. *Positive Mental Health Surveillance Indicator Framework*. Ontario, Canada; 2016.

BC Mental Health and Substance Use Services. *Health Compass Transformative Practices, Embracing Mental Wellbeing. Resource Guide for Health Care Providers* [Internet]. 2016.

Public Health Agency of Canada. *Mental Health Promotion* [Internet]. Ontario, Canada; 2014.

World Health Organization (WHO). *Mental health: a state of wellbeing* [Internet]. 2014.

World Health Organization (WHO). *Risks to Mental Health: An Overview of Vulnerabilities and Risk Factors* [Internet]. 2012.

Mental Health Protective Factors

Best Start Resource Centre. *Creating Circles of Support for Pregnant Women and New Parents: A manual for service providers supporting women's mental health in pregnancy and postpartum*. Toronto, Ontario; 2009.

Centre for Addiction and Mental Health. *What are the potential protective factors against mental health problems?* [Internet]. Toronto; 2012 [cited 2016 Jun 24].

Misri S, Kostaras X, Fox D. Partner support reduced depressive symptoms in postpartum depression. *Evid Based Ment Heal*. 2001; 4(51).

Pilkington P, Milne L, Cairns K, Whelan T. Enhancing reciprocal partner support to prevent perinatal depression and anxiety: a Delphi consensus study. *BMC Psychiatry*. 2016; 16(23).

Postpartum Blues, Postpartum Depression, Postpartum Anxiety, Postpartum Psychosis

Austin M, Highet N, Group EW. *Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline*. Melbourne, Australia; 2017.

Austin M, Highet N, Guidelines Expert Advisory Committee. *Clinical practice guidelines for depression and related disorders – anxiety, bipolar disorder and puerperal psychosis – in the perinatal period. A guideline for primary care health professionals*. [Internet]. Melbourne, Australia; 2011.

Austin M, Tully L, Parker G. Examining the relationship between antenatal anxiety and postnatal depression. *J Affect Disord*. 2007; 101(1-3):169–74.

BC Reproductive Mental Health Program & Perinatal Services. *Best Practice Guidelines for Mental Health Disorders in the Perinatal Period*. [Internet]. 2014.

Centre for Addiction and Mental Health (CAMH). *Postpartum Depression: A Guide for Front-Line Health and Social Service Providers. Clinical Overview* [Internet]. Toronto, Ontario.

Centre for Chronic Disease Prevention Public Agency of Canada. *Positive Mental Health Surveillance Indicator Framework*. Ontario, Canada; 2016.

Highet N, Stevenson A, Purtell C. Qualitative insights into women's personal experiences of perinatal depression and anxiety. *Women Birth*. 2014; 27(3):179–84.

Milgrom J, Gemmill A, Bilszta J, Hayes B, Barnett B, Brooks J, et al. Antenatal risk factors for postnatal depression: a large prospective study. *J Affect Disord*. 2008; 108(1-2):147–57.

Moore D, Ayers S, Drey N. A Thematic Analysis of Stigma and Disclosure for Perinatal Depression on an Online Forum. *JMIR Ment Heal*. 2016; 3(2).

National Institute for Health Care Management. *Identifying and treating maternal depression: strategies and considerations for health plans*. [Internet]. Washington; 2010.

National Collaborating Centre for Mental Health Royal College of Psychiatrists' Research and Training. *Antenatal and Postnatal Mental Health: The NICE guideline on Clinical Management and Service Guidance*. Updated edition. Leicester; 2014

O'Hara M, Wisner K, Asher H. Perinatal mental illness: Definition, description and aetiology. *Best Pract Res Clin Obstet Gynaecol*. 2014; 28:3–12.

Robertson E, Jones I, Haque S, Holder R CN. Risk of puerperal and non-puerperal recurrence of illness following bipolar affective puerperal (post-partum) psychosis. *Br J Psychiatry*. 2005; 186:258–9.

(SIGN) Network SIG, Scottish Intercollegiate Guidelines Network (SIGN). *Management of perinatal mood disorders* [Internet]. Edinburgh; 2012.

World Health Organization (WHO). Risks to Mental Health: An Overview of Vulnerabilities and Risk Factors [Internet]. 2012.

Postpartum Depression Screening and the Edinburgh Postnatal Depression Scale

Austin M, Hight N, Group EW. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia; 2017.

Austin M, Hight N, Guidelines Expert Advisory Committee. Clinical practice guidelines for depression and related disorders – anxiety, bipolar disorder and puerperal psychosis – in the perinatal period. A guideline for primary care health professionals. [Internet]. Melbourne, Australia; 2011.

Australian Population Health Development Principal, Screening Subcommittee. Population Based Screening Framework [Internet]. 2008.

BC Reproductive Mental Health Program & Perinatal Services. Best Practice Guidelines for Mental Health Disorders in the Perinatal Period. [Internet]. 2014.

Bowen A, Bowen R, Butt P, Rahman K, Muhajarine N. Patterns of depression and treatment in pregnant and postpartum women. *Can J Psychiatry*. 2012; 57(3):161–7.

Cox J, Holden J. Perinatal Mental Health: A guide to the Edinburgh Postnatal Depression Scale. London: Gaskell; 2003.

Cox J, Holden J, Henshaw C. Perinatal Mental Health: The Edinburgh Postnatal Depression Scale (EPDS) Manual. 2nd ed. RC Psych Publications; 2014.

Department of Health. Edinburgh Postnatal Depression Scale (EPDS): Translated versions – validated. Perth, Western Australia; 2006.

Gibson J, McKenzie-McHarg K, Shakespeare J, Price J, Gray R. A systematic review of studies validating the Edinburgh Postnatal Depression Scale in antepartum and postpartum women. *Acta Psychiatr Scand*. 2009; 119(5):350–64.

International Epidemiological Association. A Dictionary of Epidemiology [Internet]. 6th ed. Porta M, editor. New York, NY: Oxford University Press; 2014.

Kingston D, McDonald S, Tough S, Austin M-P, Hegadoren K, Lasiuk G. Public views of acceptability of perinatal mental health screening and treatment preference: a population based survey. *BMC Pregnancy Childbirth*. 2014 Dec; 14(1):67.

Kroenke K, Spitzer R, Williams J. The Patient Health Questionnaire-2: Validity of a Two-Item Depression Screen. *Med Care* [Internet]. 2003; 41:1284–94

Small R, Lumley J, Yelland J, Brown S. The performance of the Edinburgh Postnatal Depression Scale in English speaking and non-English speaking populations in Australia. *Soc Psychiatry Psychiatr Epidemiol*. 2007; 42(1):70–80.

Moore D, Ayers S, Drey N. A Thematic Analysis of Stigma and Disclosure for Perinatal Depression on an Online Forum. *JMIR Ment Heal*. 2016; 3(2).

National Collaborating Centre for Mental Health Royal College of Psychiatrists' Research and Training. Antenatal and Postnatal Mental Health: The NICE guideline on Clinical Management and Service Guidance. Updated edition. Leicester; 2014

(SIGN) Network SIG, Scottish Intercollegiate Guidelines Network (SIGN). Management of perinatal mood disorders [Internet]. Edinburgh; 2012.

US Commission on Chronic Illness. Chronic Illness in the US. Vol.1. Prevention of Chronic Illness. Cambridge, Mass; 1957.

Postpartum Depression Treatment and Self Care

Austin M, Highet N, Group EW. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia; 2017.

Austin M, Highet N, Guidelines Expert Advisory Committee. Clinical practice guidelines for depression and related disorders – anxiety, bipolar disorder and puerperal psychosis – in the perinatal period. A guideline for primary care health professionals. [Internet]. Melbourne, Australia; 2011.

Alberta Health Services. Healthy Parents, Healthy Children [Internet]. 2014.

BC Reproductive Mental Health Program & Perinatal Services. Best Practice Guidelines for Mental Health Disorders in the Perinatal Period. [Internet]. 2014.

Canadian Society of Exercise Physiology. PAR med-X for PREGNANCY Physical Activity Readiness Medical Examination [Internet]. 2016.

Kingston D, McDonald S, Tough S, Austin M-P, Hegadoren K, Lasiuk G. Public views of acceptability of perinatal mental health screening and treatment preference: a population based survey. *BMC Pregnancy Childbirth*. 2014 Dec; 14(1):67.

Liberto T. Screening for depression and help-seeking in postpartum women during well-baby pediatric visits: an integrated review. *J Pediatr Heal Care* [Internet]. 2012; 26(2):109–17.

National Collaborating Centre for Mental Health Royal College of Psychiatrists' Research and Training. Antenatal and Postnatal Mental Health: The NICE guideline on Clinical Management and Service Guidance. Updated edition. Leicester; 2014.

National Collaborating Centre for Mental Health Royal College of Psychiatrists' Research and Training. Antenatal and Postnatal Mental Health: The NICE Guideline on Clinical Management and Service Guidance. Leicester; 2007.

(SIGN) Network SIG, Scottish Intercollegiate Guidelines Network (SIGN). Management of perinatal mood disorders [Internet]. Edinburgh; 2012.

Canadian Society of Exercise Physiology. PAR med-X for PREGNANCY Physical Activity Readiness Medical Examination [Internet]. 2016.

Prevalence: PPD and Mental Illness

Anxiety and Depression Association of America. Understanding anxiety and depression. 2016.

Austin M. Antenatal screening and early intervention for perinatal distress depression and anxiety: where to from here? *Arch Women's Ment Heal*. 2004; 7:1–6.

Banti S, Mauri M, Oppo A. From the third month of pregnancy to 1 year postpartum. Prevalence, incidence, recurrence, and new onset of depression. Results from the perinatal Depression-Research & Screening study. *Compr Psychiatry* [Internet]. 2011; 52(4):343–51.

Canadian Mental Health Association Alberta. Mental Illness in Canada.

Coates A, Schaefer C, JL A. Detection of postpartum depression and anxiety in a large health plan. *J Behav Heal Serv Res*. 2004; 31(2):117–33.

Gaynes B, Gavin S, Meltzer-Brody S, Lohr K, Swinson T, Gartlehner G, et al. Perinatal depression: prevalence, screening accuracy, and screening outcomes [Internet]. Rockville,MD; 2005.

Gavin NI, Gaynes BN, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T. Perinatal depression: A systematic review of prevalence and incidence. *Obstet Gynecol*. 2005; 106(5):1071–83.

Grace S, Evindar A, Stewart D. The effect of postpartum depression on child cognitive development and behavior: a review and critical analysis of the literature. *Arch Womens Mentl Heal*. 2003; 6(4):263–74

Horwitz S, Briggs-Gowan M, Storfer-Isser A, Carter A. Persistence of maternal depressive symptoms throughout the early years of childhood. *J Womens Heal*. 2009; 18(5):637–45.

Kingston D, Heaman M, Fell D, Dzakpasu S, Chalmers B. Factors associated with perceived stress and stressful life events in pregnant women: findings from the Canadian Maternity Experiences Survey. *Matern Child Heal J*. 2012; 16(1):158–68.

Liberto T. Screening for depression and help-seeking in postpartum women during well-baby pediatric visits: an integrated review. *J Pediatr Heal Care* [Internet]. 2012; 26(2):109–17.

Milgrom J, Gemmill A, Bilszta J, Hayes B, Barnett B, Brooks J, et al. Antenatal risk factors for postnatal depression: a large prospective study. *J Affect Disord*. 2008; 108(1-2):147–57.

Potestio M. Treated Postpartum Depression Prevalence in Alberta. Unpublished raw data. Alberta Health Services; Research and Innovation: Surveillance and Infrastructure, Population Public and Aboriginal Health. 2012.

Priest S, Austin M, Barnett B, Buist A. A psychosocial risk assessment model (PRAM) for use with pregnant and postpartum women in primary care settings. *Arch Womens Ment Heal* 2008. 2008; 11(5-6):307–17.

Spitzer R, Williams J, Kroenke K. Validity and utility of the PRIME-MD patient health questionnaire in assessment of 3000 obstetric-gynecologic patients: the PRIME-MD Patient Health Questionnaire Obstetrics-Gynecology Study. *Am J Obs Gynecol*. 2000; 183(3):759–69.

Underwood L, Waldie K, D'Souza S, Peterson E, Morton S. A review of longitudinal studies on antenatal and postnatal depression. *Arch Women's Ment Heal*. 2016; 19:711–20.

Suicide Risk

American Psychiatric Association. Practice Guidelines for the Psychiatric Evaluation of Adults. [Internet]. 3rd ed. Arlington, VA; 2016.

Austin M, Hight N, Group EW. Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline. Melbourne, Australia; 2017.

BC Reproductive Mental Health Program & Perinatal Services. Best Practice Guidelines for Mental Health Disorders in the Perinatal Period. [Internet]. 2014.

Banti S, Mauri M, Oppo A. From the third month of pregnancy to 1 year postpartum. Prevalence, incidence, recurrence, and new onset of depression. Results from the perinatal Depression-Research & Screening study. *Compr Psychiatry* [Internet]. 2011; 52(4):343–51.

Borges G, Nock M, Haro Abad J. Twelve-month prevalence of and risk factors for suicide attempts in the World Health Organization World Mental Health Surveys. *J Clin Psychiatry*. 2010; 71(12):1617–28.

Centre for Addiction and Mental Health. CAMH Suicide Prevention and Assessment Handbook [Internet]. 2011. 8-9 p.

Centre for Addiction and Mental Health. What are the potential protective factors against mental health problems? [Internet]. Toronto; 2012 [cited 2016 Jun 24].

Centers for Disease Control and Prevention. Suicide: Risk and Protective Factors [Internet]. [Cited 2016 Jun 24].

Cox J, Holden J, Henshaw C. Perinatal Mental Health: The Edinburgh Postnatal Depression Scale (EPDS) Manual. 2nd ed. RC Psych Publications; 2014

Department of Veterans Affairs. Assessment and Management of Risk for Suicide Working Group: VA/DoD Clinical Practice Guideline for Assessment and Management of Patients at Risk for Suicide. [Internet]. Washington, DC; 2013

Hight N, Stevenson A, Purtell C. Qualitative insights into women's personal experiences of perinatal depression and anxiety. *Women Birth*. 2014; 27(3):179–84

Kingston D, McDonald S, Tough S, Austin M-P, Hegadoren K, Lasiuk G. Public views of acceptability of perinatal mental health screening and treatment preference: a population based survey. *BMC Pregnancy Childbirth*. 2014 Dec; 14(1):67.

Lindahl V, Pearson J, Colpe L. Prevalence of suicidality during pregnancy and the postpartum. *Arch Women's Ment Heal*. 2005; 8:77–87.

Moore D, Ayers S, Drey N. A Thematic Analysis of Stigma and Disclosure for Perinatal Depression on an Online Forum. *JMIR Ment Heal*. 2016; 3(2).

O'Hara M, Wisner K, Asher H. Perinatal mental illness: Definition, description and aetiology. *Best Pract Res Clin Obstet Gynaecol*. 2014; 28:3–12.

Sokero P, Melartin T, Rytsala H, Leskela U, Lestela-Mielonen P, Istometsa E. Prospective study of risk factors for attempted suicide among patients with DSM IV major depressive disorder. *Br J Psychiatry*. 2005; 186:314–8.

Tidemalm D, Langstrom N, Lichtenstein P, Runeson B. Risk of suicide after suicide attempt according to coexisting psychiatric disorder: Swedish cohort study with long term follow-up. *BMJ*. 2008; 337(a2205).

Turecki G, Brent D. Suicide and suicidal behaviour. *Lancet*. 2016; 387(10024):1227–39.

Wisner K, Sit D, McShea M, Rizzo D, Zoretich R. Onset Timing, Thoughts of Self-harm, and Diagnoses in Postpartum Women With Screen-Positive Depression Findings. *JAMA Psychiatry* [Internet]. 2013; 70(5):490–8.

Values Beliefs and Attitudes and Patient and Family Centred Care

Centre for Addiction and Mental Health. A Review of Clinical Cultural Competence: Definitions, Key Components, Standards and Selected Trainings. 2004; 19–41.

College and Association of Registered Nurses of Alberta. Practice Standards for Regulated Members with the Canadian Nurses Association Code of Ethics for Registered Nurses. Edmonton; 2008.

Institute for Patient and Family-Centred Care. Institute for Patient and Family-Centred Care. Frequently Asked Questions. [Internet]. 2010 [cited 2016 Feb 20].

Megnin-Viggars O, Symington I, Howard L. Experience of care for mental health problems in the antenatal or postnatal period for women in the UK: a systematic review and meta-synthesis of qualitative research. *Arch Womens Ment Heal*. 2015; 18(6):745–59.

Quality and Healthcare Improvement Engagement and Patient Experience. Patient and Family Centred Care Resource Kit [Internet]. Calgary, Alberta; 2014.

Yiu V, Gordon D, Woods S, Pougnet J. The Patient First Strategy [Internet]. Calgary, Alberta; 2015.