

Post COVID Rehabilitation Continuing Care – Frequency of Screening, Assessment and Reassessment

Rehabilitation needs are identified for adults 18 years and older during:

- Screening – identifies any new symptom, change in condition or reduction in function after recovery from COVID-19.
- Care plan review – identifies any new symptom, change in condition or reduction in function after recovery from COVID-19.

Admission to Continuing Care Programs

- Upon admission to a continuing care program, complete or review post COVID-19 screening based on a previous diagnosis of COVID-19 and the applicable client group, clinical area or program.
- Complete comprehensive admission assessment following established clinical area or program timelines.
- Discuss post COVID-19 symptoms and any recommended interventions with patient, family or decision-maker and interdisciplinary team for care and treatment plan development.
- Provide education to the patient, family or decision maker regarding risk factors and strategies for management of long-term recovery from post COVID-19 symptoms.
- Encourage self-management of rehabilitation needs (as able or with the support of family or caregiver).

Prioritization of referrals, assessment and reassessment

- Referrals, assessment and reassessment of patients with post COVID-19 symptoms should be prioritized following routine referral and treatment processes.
- Prioritization should include consideration of risk for acute care admission, risk for higher level of care needs, patient struggling to manage in current care stream, risk of social isolation, length of outbreak (longer outbreak = higher risk for functional, cognitive and overall decline).

Discharge from Continuing Care Programs

- Provide discharge instructions and education to patient and family or caregiver.
- Ensure patient and family or caregiver have required information to initiate and maintain self-management of rehabilitation needs.
- Communicate discharge and any follow-up needs to primary care provider (Primary Care Network, physician, nurse practitioner).

For more information
continuingcare@ahs.ca



Policy, Practice, Access and Case Management
 Provincial Seniors Health and Continuing Care

Patient Population, Clinical Area or Program	Initial Screening	Assessment	Reassessment
Post-Acute: Adult 18 years and older			
Sub-Acute Restorative Care	<ul style="list-style-type: none"> • Review screening completed in acute care within 24 hours of admission, (ideally within 8 hours) OR if no screening has been completed: <ul style="list-style-type: none"> ○ Complete within 7 days after recovery (14 days from positive test or absence of symptoms) and/or ○ Complete within 24 hours before discharge or ○ Complete within 24 hours before transfer to another unit and/or facility 	<ul style="list-style-type: none"> • Complete comprehensive assessment and treatment plan following established timeframes • Identify Allied Health disciplines needing to be included in interdisciplinary team (IDT) • Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> • Evaluate and reassess treatment plan at 3, 6 and 12-month intervals and when there is a significant change in patient's health status or functional ability • Reassess before transfer to another unit and/or facility
Home Care Home Living and Non-Designated Supportive Living (Adult 18 years and older)			
Short Term Acute Client Group (STAC)	<ul style="list-style-type: none"> • Review screening completed in acute care during established intake process and upon first visit, or • Complete the appropriate screening tool with established intake process or upon first visit 	<ul style="list-style-type: none"> • Complete comprehensive assessment and treatment plan following established timeframes • Identify Allied Health disciplines needing to be included in IDT • Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> • When post COVID-19 rehabilitative need is identified, reassess before discharge to establish self-management plan • Evaluate effectiveness of rehabilitation intervention plan

Policy, Practice, Access and Case Management
 Provincial Seniors Health and Continuing Care

Patient Population, Clinical Area or Program	Initial Screening	Assessment	Reassessment
Acute, Rehab, End of Life and Wellness Client Groups	<ul style="list-style-type: none"> Review screening completed in acute care during established intake process and upon first visit, or Complete the appropriate screening tool with established intake process or upon first visit 	<ul style="list-style-type: none"> Complete comprehensive assessment and treatment plan following established timeframes Identify Allied Health disciplines needing to be included in IDT Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> When post COVID-19 rehabilitative need is identified, reassess at 3, 6 and 12-month intervals and/or before discharge to establish self-management plan (as appropriate) Reassess when there is a significant change in health status (e.g. change in functional status after acute care admission or due to symptom exacerbation) Evaluate effectiveness of rehabilitation intervention plan
Long Term Supportive/Maintenance Client Groups	<ul style="list-style-type: none"> Review screening completed in acute care during established intake process and upon first visit, or Complete the appropriate screening tool with established intake process or upon first visit 	<ul style="list-style-type: none"> Complete comprehensive assessment and treatment plan following established timelines Identify Allied Health disciplines needing to be included in IDT Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> When post COVID-19 rehabilitative need hasn't previously been identified, screen for change in status at 3, 6 and 12-month intervals When post COVID-19 rehabilitative need has been identified, reassess treatment and intervention plan at 3, 6 and 12-month intervals, and Reassess when there is a significant change in health status (see interRAI definition)

Policy, Practice, Access and Case Management
 Provincial Seniors Health and Continuing Care

Patient Population, Clinical Area or Program	Initial Screening	Assessment	Reassessment
			for significant change in status) • Evaluate effectiveness of rehabilitation intervention plan
Designated Supportive Living (Adult 18 years and older)			
Long Term Supportive/Maintenance Client Groups	<ul style="list-style-type: none"> Review screening completed in acute care within 24 hours of admission, ideally within 8 hours or Complete the appropriate screening tool with established intake process or within 24 hours of admission 	<ul style="list-style-type: none"> Complete comprehensive assessment and treatment plan following established timelines Identify Allied Health disciplines needing to be included in IDT Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> When no prior post COVID-19 rehabilitative need is identified, screen for change in status at 3, 6 and 12-month intervals When post COVID-19 rehabilitative need is identified, reassess treatment and intervention plan at 3, 6 and 12-month intervals, and Reassess when there is a significant change in health status (see interRAI definition for significant change in status) Evaluate effectiveness of rehabilitation intervention plan

Policy, Practice, Access and Case Management
 Provincial Seniors Health and Continuing Care

Patient Population, Clinical Area or Program	Initial Screening	Assessment	Reassessment
Long Term Care (LTC) Facility Living (Adult 18 years and older)			
Long Term Care (LTC)	<ul style="list-style-type: none"> Review screening completed in acute care within 24 hours of admission, ideally within 8 hours or Complete the appropriate screening tool with established intake process or within 24 hours of admission 	<ul style="list-style-type: none"> Complete comprehensive assessment and treatment plan following established timelines Identify Allied Health disciplines needing to be included in IDT Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> When no prior post COVID-19 rehabilitative need is identified, screen for change in status at 3, 6 and 12-month intervals When post COVID-19 rehabilitative need is identified, reassess treatment and intervention plan quarterly, and Reassess when there is a significant change in health status (see interRAI definition of significant change in status) Evaluate effectiveness of rehabilitation intervention plan
Hospice (Adult 18 years and older)			
Hospice	<ul style="list-style-type: none"> Review screening completed in acute care within 24 hours of admission, ideally within 8 hours or Complete the appropriate screening tool with established intake process or within 24 hours of admission 	<ul style="list-style-type: none"> Complete comprehensive assessment and treatment plan following established timelines Identify Allied Health disciplines needing to be included in IDT Initiate rehabilitation intervention plan 	<ul style="list-style-type: none"> When post COVID-19 rehabilitative need is identified, reassess treatment and intervention plan following clinical need related to symptom management, and Reassess with significant change in health status (e.g. change in functional status) Evaluate effectiveness of rehabilitation intervention plan.