



Document Number: GLS.210

Revision Number: 2.43

Document Type: Procedure

Effective Date: 12/7/2022 1:32:59 PM

Location: 1. AHS Folder Structure\Document Administration\Guide to Lab Services\Information for Lab and Health Professionals\Reference Ranges\Central Zone

GLS.210 – DTH Meditech HCIS Reference Intervals

Applicability This document applies to Central Zone Red Deer EPIC HCIS personnel of AHS Laboratory Services.

Purpose This document states the reference intervals for Central Zone Red Deer Chemistry EPIC HCIS Laboratory analytes.

Test	Reference Intervals																												
ACETAMINOPHEN, P/S	No reference interval																												
ALBUMIN/CREATININE RATIO, U	MALE AND FEMALE mg/mmol <1 m <17.50 1 m – <2 y <4.00 2 – 150 y <3.00																												
ALBUMIN, P/S	MALE, FEMALE, U/X g/L 0 – 364 d 22 – 45 365 d – 150 y 30 – 45																												
ALBUMIN, TIMED, 24 Hour Urine	Urine Albumin Excretion Rate MALE AND FEMALE ug/min 0 – 150 y <20																												
ALKALINE PHOSPHATASE (ALP), P/S	<table border="0"> <thead> <tr> <th></th> <th>Male U/L</th> <th>Female</th> <th>X/U</th> </tr> </thead> <tbody> <tr> <td>Age < 15 d</td> <td>70 – 320</td> <td>70 – 320</td> <td>70 - 320</td> </tr> <tr> <td>15 d to < 1 y</td> <td>130 – 500</td> <td>130 – 500</td> <td>130 - 500</td> </tr> <tr> <td>1 to < 13 y</td> <td>130 - 430</td> <td>130 – 430</td> <td>130 - 430</td> </tr> <tr> <td>13 to < 15 y</td> <td>130 - 500</td> <td>60 – 225</td> <td>60 - 500</td> </tr> <tr> <td>15 to < 18 y</td> <td>60 – 250</td> <td>50 – 140</td> <td>50 – 250</td> </tr> <tr> <td>18 to 150 y</td> <td>40 – 120</td> <td>40 – 120</td> <td>40 - 120</td> </tr> </tbody> </table>		Male U/L	Female	X/U	Age < 15 d	70 – 320	70 – 320	70 - 320	15 d to < 1 y	130 – 500	130 – 500	130 - 500	1 to < 13 y	130 - 430	130 – 430	130 - 430	13 to < 15 y	130 - 500	60 – 225	60 - 500	15 to < 18 y	60 – 250	50 – 140	50 – 250	18 to 150 y	40 – 120	40 – 120	40 - 120
	Male U/L	Female	X/U																										
Age < 15 d	70 – 320	70 – 320	70 - 320																										
15 d to < 1 y	130 – 500	130 – 500	130 - 500																										
1 to < 13 y	130 - 430	130 – 430	130 - 430																										
13 to < 15 y	130 - 500	60 – 225	60 - 500																										
15 to < 18 y	60 – 250	50 – 140	50 – 250																										
18 to 150 y	40 – 120	40 – 120	40 - 120																										
ALANINE AMINOTRANSFERASE (ALT), P/S	M,F,U/X <18 y <40 U/L M ≥ 18 y <70 U/L F ≥ 18 y <50 U/L U/X ≥ 18 y <50 U/L																												
AMMONIA, P	MALE AND FEMALE µmol/L 0 – <3 m 30-100 3m – 150 y 20-50																												
ASPARTATE AMINOTRANSFERASE (AST), P/S	MALE AND FEMALE U/X U/L <1 m <115 1m to <1 y <80 1y to <7 y <60 7y to < 18 y <45 18-150 y M U/X <55 18-150 y F <45																												

Test	Reference Intervals
BETA HUMAN CHORIONIC GONADOTROPIN (BCHG), Urine	Negative
BETA HUMAN CHORIONIC GONADOTROPIN (BHCG), P/S	MALE AND FEMALE IU/L 0 – 150 y < 5
BETA-HYDROXYBUTYRATE, B	MALE AND FEMALE mmol/L 0 - 150 y <0.4
BILIRUBIN – DIRECT, P/S	MALE, FEMALE and U/X µmol/L 0 – 150 y < 7
BILIRUBIN – TOTAL, P/S	MALE, FEMALE and U/X µmol/L 30d to 150 y <20
BILIRUBIN- Neonatal	M F U/X umol/L <7d no range 7 to 15d <250 15 d-<30d <20
BLOOD GASES	Refer to tables on the last page of this document
BNP (B-TYPE NATRIURETIC PEPTIDE), P	MALE AND FEMALE ng/L 0 – 150 y Diagnostic Criteria: <100 CHF Unlikely 100 – 500 CHF Possible >500 CHF Likely
NT -pro BNP, P/S	MALE FEMALE ng/L Age < 1 y 54-556 1y to < 2 y 39-578 2y to < 6 y 20-565 6y to < 12 y 10-340 12y to-< 18 y 6-216 18y to 150 0-300
CALCIUM, 24 Hour Urine	MALE AND FEMALE mmol/d 0 – 150 y 2.50-7.50
CALCIUM, P/S	MALE, FEMALE and UNKNOWN mmol/L 0 – 10 d 1.80 – 2.90 11 – 365 d 2.20 – 2.80 > 1y 2.10 – 2.60
CALCIUM IONIZED, POST FILTER	MALE AND FEMALE mmol/L 0-18 yr 0.35 - 0.45 18 – 150 yr 0.20 – 0.40
CALCIUM, IONIZED, S	MALE AND FEMALE mmol/L 0 – 29 days 0.90 – 1.30 29 days – 150 yr 1.15 – 1.35
CARBAMAZEPINE, P/S	MALE AND FEMALE µmol/L Therapeutic Range 0 – 150 y 17 – 50
CBC (COMPLETE BLOOD COUNT), B	See "DTH Meditech HCIS CBC and Differential Reference Intervals"
CELL COUNT, CSF	Color Colorless Appearance Clear WBC 0 – 5 x10 ⁶ /L

Test	Reference Intervals
	RBC 0 x 10 ⁶ /L
CELL COUNT, Fluid	Transudates Color Colorless Clarity Clear RBC < 10,000 x 10 ⁶ /L WBC < 1,000 x 10 ⁶ /L Exudates Color Variable Clarity Turbid RBC > 10,000 x 10 ⁶ /L WBC > 1,000 x 10 ⁶ /L
CHLAMYDIA/GONORRHOEAE PROBE NUCLEIC ACID AMPLIFICATION TEST (NAAT)	Negative
CHOLESTEROL, TOTAL, P/S	MALE AND FEMALE: mmol/L <2 y 2.36 – 5.32 2 – <18 y 2.70 – 5.89 The following comment will be attached to the result: Acceptable limit relative to dyslipidemia and atherosclerosis risk is < 4.40 mmol/L. 18 – 150 y No reference range The following comment will be attached to the result: Desirable < 5.17 mmol/L High >= 6.21 mmol/L
CLOSTRIDIUM DIFFICILE	Negative
CO-OXIMETRY	Refer to tables on the last page of this document
CREATINE KINASE (CK), P/S	MALE U/X U/L 0-150y 30-350 FEMALE U/L 0-150 y 30-200
C-REACTIVE PROTEIN (CRP), P/S	MALE AND FEMALE mg/L 0 – 150 y <8.1
CREATININE, 24 Hour Urine	MALE FEMALE U/X mmol/d Age < 3 y no range 3 to < 8 y 1.0-6.0 9 to < 12 y 5.0-12.5 13 to < 17 y 7.0-16.5 18 to 150 y M 9.0-18.0 18 to 150 y F 7.0-16.0 18 to 150 y U/X 7.0-18.0
CREATININE, P/S	MALE, FEMALE, U/X umol/L <1d no range 1d to <2 y 10 – 40 2y to <6 y 20 – 45

Test	Reference Intervals
	6y to <13 y 20 – 75 13y to <15 y 30 – 95 15-150 y M 50-120 15-150 y F 40-100 15-150 y U/X 40-120
CREATININE CLEARANCE, 24 Hour Urine + P/S	MALE AND FEMALE mL/min/1.73 m ² 78.00-138.00
CRYOGLOBULIN, S	Negative
D-Dimer, Quantitative, P	0.00 - 0.50 mg/L FEU A D-Dimer BELOW the 0.50 mg/L FEU cutoff may be used with a standardized Clinical Assessment and/or imaging studies to help exclude venous thromboembolism (VTE). Values above the cutoff are not diagnostically useful in VTE assessment.
D-DIMER QUANTITATIVE (Triage Meter), B	0.5 mg/L A D-Dimer level BELOW the 0.5 mg/L cutoff may be used with the standardized clinical assessment and/or imaging studies to help exclude venous thromboembolism (VTE). Values above the cutoff are not diagnostically useful in VTE assessment.
DIFFERENTIAL, B	See "DTH Meditech HCIS CBC and Differential Reference Intervals".
DIGOXIN, P/S	MALE AND FEMALE nmol/L 0 – 150 y Suggested Ranges: Heart failure 0.6 – 1.2 Atrial Fibrillation Not Defined Caution: Results > 1.5 nmol/L are associated with a higher risk of toxicity in heart failure patients.
DNA DOUBLE STRAND ANTIBODY, S	Negative
eGFR (CKD-EPI)	MALE FEMALE U/X 18 -150 y >59 mL/min/1.73m ² The following Interpretive Comment will be appended to all eGFR results on adults (≥18 years) with eGFR results <60 mL/min/1.73 m ² . "eGFR <60 mL/min/1.73 m ² or urine Albumin/creatinine ratio ≥3.00 mg/mmol for more than 3 months suggests chronic kidney disease. For information on diagnosis, management and referral see www.diagnoseckd.ca ." The following Interpretive Comment will be appended to all eGFR results on adults (≥18 years) "Reduced muscle mass will lead to overestimation, and increased muscle mass underestimation of eGFR."

Test	Reference Intervals
	<p>eGFR results will not be reported on patients <18 years of age or dialysis patients.</p> <p>The following comment will be added to all eGFR results on patients of unknown gender.</p> <p>“Unable to calculate as the gender is unknown.”</p>
ELECTROLYTES, P/S	<p>MALE, FEMALE, U/X mmol/L</p> <p>Sodium 0 – 150 y 135 – 145</p> <p>Potassium 0 – <29 d 3.5 – 6.0 29d – <1 y 3.5 – 5.5 1y – <18 y 3.5 – 5.0 18-150 y 3.5 – 5.0</p> <p>NOTE: Potassium concentration is 0.2 to 0.5 mmol/L higher in serum (gold top).</p> <p>Chloride 0 – 150 y 98- 112</p> <p>CO2 0 – 150 y 20 -s32</p>
ELECTROLYTES, 24 Hour Urine	<p>MALE AND FEMALE mmol/d Intake dependent</p> <p>Sodium 0-150 y 45-250</p> <p>Potassium 0 – 150 y 25- 125</p> <p>Chloride 0-150 y 110-250</p>
ESR (ERYTHROCYTE SEDIMENTATION RATE), B	<p>0 – 17 y 0 – 10 mm/h</p> <p>FEMALE 18 – 150 y 0 – 20 mm/h</p> <p>MALE 18 – 150 Y 0 – 15 mm/h</p>
ETHANOL, P/S	<p>MALE AND FEMALE mmol/L 0 – 150 y <2</p>
FERRITIN, P/S	<p>MALE, FEMALE U/X µg/L</p> <p>0 – <6 m 50-500 6m– <16 y 15-100</p>

Test	Reference Intervals																						
	<p>MALE, 16-150 y 30-500</p> <p>FEMALE 16-150 y 20-300</p> <p>U/X 16-150 y 20-500</p>																						
FIBRINOGEN, QUANTITATIVE, P	1.62 – 4.24 g/L																						
FSH (FOLLICLE STIMULATING HORMONE), P/S	<p>IU/L</p> <table border="1"> <thead> <tr> <th colspan="2">Female and Unknown</th> </tr> <tr> <th>Age</th> <th>Interval</th> </tr> </thead> <tbody> <tr> <td><2y</td> <td><8.0</td> </tr> <tr> <td>2-5y</td> <td><5.0</td> </tr> <tr> <td>6-10y</td> <td><4.0</td> </tr> <tr> <td>11-150y</td> <td>See comment Follicular: 2.0-10.0 Luteal: 1.0-9.0 Midcycle: 3.0-33.0 Post menopausal: 23.0-116.0</td> </tr> <tr> <th colspan="2">Male</th> </tr> <tr> <th>Age</th> <th>Interval</th> </tr> <tr> <td><11y</td> <td><3.0</td> </tr> <tr> <td>11-12y</td> <td><9.0</td> </tr> <tr> <td>13-150y</td> <td>1.0-18.0</td> </tr> </tbody> </table>	Female and Unknown		Age	Interval	<2y	<8.0	2-5y	<5.0	6-10y	<4.0	11-150y	See comment Follicular: 2.0-10.0 Luteal: 1.0-9.0 Midcycle: 3.0-33.0 Post menopausal: 23.0-116.0	Male		Age	Interval	<11y	<3.0	11-12y	<9.0	13-150y	1.0-18.0
Female and Unknown																							
Age	Interval																						
<2y	<8.0																						
2-5y	<5.0																						
6-10y	<4.0																						
11-150y	See comment Follicular: 2.0-10.0 Luteal: 1.0-9.0 Midcycle: 3.0-33.0 Post menopausal: 23.0-116.0																						
Male																							
Age	Interval																						
<11y	<3.0																						
11-12y	<9.0																						
13-150y	1.0-18.0																						
GAMMA-GLUTAMYL TRANSFERASE (GGT), P/S	<p>MALE, FEMALE, U/X U/L</p> <p>0 – <15 d 20 – 200</p> <p>15 d – 1 y <100</p> <p>1yr - <18 y <27</p> <p>MALE U/X</p> <p>18 – 150 y <80</p> <p>FEMALE</p> <p>18 – 150 Y <50</p>																						
GENTAMICIN – 8 hour INTERVAL, P/S GENTAMICIN – post, P/S GENTAMICIN – OTHER, P/S GENTAMICIN – PRE, P/S	<table border="1"> <thead> <tr> <th colspan="2">Therapeutic Target for Multiple daily dosing</th> </tr> </thead> <tbody> <tr> <td>Pre -dose</td> <td><2.0 mg/L</td> </tr> <tr> <td>Post- dose</td> <td>5.0-1.0 mg/L</td> </tr> <tr> <th colspan="2">Therapeutic target for extended Interval dosing</th> </tr> <tr> <td>Pre-dose</td> <td>0.0-1.0 mg/L</td> </tr> <tr> <td>Post-dose</td> <td>Contraindicated for adults with normal renal function.</td> </tr> </tbody> </table>	Therapeutic Target for Multiple daily dosing		Pre -dose	<2.0 mg/L	Post- dose	5.0-1.0 mg/L	Therapeutic target for extended Interval dosing		Pre-dose	0.0-1.0 mg/L	Post-dose	Contraindicated for adults with normal renal function.										
Therapeutic Target for Multiple daily dosing																							
Pre -dose	<2.0 mg/L																						
Post- dose	5.0-1.0 mg/L																						
Therapeutic target for extended Interval dosing																							
Pre-dose	0.0-1.0 mg/L																						
Post-dose	Contraindicated for adults with normal renal function.																						
GLUCOSE, CSF	<p>MALE AND FEMALE mmol/L</p> <p>0 - 150 y 2.2 – 4.4</p>																						
GLUCOSE – FASTING, P/S	<p>MALE AND FEMALE mmol/L</p> <p>< 30 d 2.5 – 5.5</p> <p>30 d – 150 y 3.3 – 6.0</p>																						

Test	Reference Intervals
GLUCOSE – RANDOM, P/S	MALE AND FEMALE mmol/L 0 – 29 d 2.5 – 11.0 30 d – 150 y 3.3 – 11.0
GLUCOSE – GESTATIONAL DIABETIC SCREEN, P/S	Gestational Diabetic Screen Interpretive Guidelines (Canadian Diabetes Guidelines 2013): Normal: < 7.8 mmol/L Follow-up with GTT-2h Pregnant: 7.8-11.0 mmol/L Gestational Diabetes: >= 11.1 mmol/L
GLUCOSE – GESTATIONAL TOLERANCE TEST, P/S	Gestational Glucose Tolerance Test Interpretive Guidelines (Canadian Diabetes Guidelines (2013)): Gestational Diabetes Mellitus - glucose results meet ONE of the criteria below: Fasting: >= 5.3 mmol/L 1h Glucose: >= 10.6 mmol/L 2h Glucose: >= 9.0 mmol/L
GLUCOSE – TOLERANCE TEST – 2 h, P/S	Glucose Tolerance Test Interpretive Guidelines (Non-pregnant; Canadian Diabetes Guidelines (2013)): Normal Glucose Tolerance: Fasting Glucose: 3.3 - 6.0 mmol/L AND 2h Glucose < 7.8 mmol/L Impaired Fasting Glucose and Impaired Glucose Tolerance: Fasting Glucose: 6.1 - 6.9 mmol/L AND 2h Glucose 7.8 - 11.0 mmol/L Diabetes Mellitus: Fasting Glucose: > 6.9 mmol/L OR 2h Glucose > 11.0 mmol/L
HIV Rapid Antibody Screen, S	MALE AND FEMALE Non-Reactive
IMMUNOGLOBULINS (QUANTITATIVE), P/S	IgG MALE AND FEMALE g/L 1d – 2 m 2.60 – 14.00 3m – 11 m 2.80 – 16.00 1 – 2 y 4.00 – 16.00 3 – 5 y 5.40 - 16.00 6 – 7 y 5.80 - 16.00 8 – 10 y 6.20 - 17.00 11 – 15 y 6.40 - 17.00 16 – 150 y 6.80 - 18.00 IgA MALE AND FEMALE g/L 1d – 2 m <1.21 3m – 5 m 0.05 – 1.20 6m – 12 m 0.10 – 1.20

Test	Reference Intervals																												
	<table> <tr><td>1 y</td><td>0.20 – 1.60</td></tr> <tr><td>2 y</td><td>0.30 – 2.00</td></tr> <tr><td>3 – 5 y</td><td>0.35 – 2.40</td></tr> <tr><td>6 – 7 y</td><td>0.40 – 2.80</td></tr> <tr><td>8 – 10 y</td><td>0.45– 3.20</td></tr> <tr><td>11 – 15 y</td><td>0.50 – 3.80</td></tr> <tr><td>16 – 150 y</td><td>0.60 – 4.20</td></tr> </table> <p><u>IgM</u> MALE AND FEMALE g/L</p> <table> <tr><td><5 m</td><td>0.14 – 1.40</td></tr> <tr><td>5 m- 3 y</td><td>0.20 – 1.60</td></tr> <tr><td>4 –6 y</td><td>0.20 – 2.10</td></tr> <tr><td>3 – 5 y</td><td>0.35 – 2.10</td></tr> <tr><td>7 – 11 y</td><td>0.30 – 2.10</td></tr> <tr><td>12 – 19 y</td><td>0.30 – 2.40</td></tr> <tr><td>20 – 150 y</td><td>0.40 – 3.00</td></tr> </table>	1 y	0.20 – 1.60	2 y	0.30 – 2.00	3 – 5 y	0.35 – 2.40	6 – 7 y	0.40 – 2.80	8 – 10 y	0.45– 3.20	11 – 15 y	0.50 – 3.80	16 – 150 y	0.60 – 4.20	<5 m	0.14 – 1.40	5 m- 3 y	0.20 – 1.60	4 –6 y	0.20 – 2.10	3 – 5 y	0.35 – 2.10	7 – 11 y	0.30 – 2.10	12 – 19 y	0.30 – 2.40	20 – 150 y	0.40 – 3.00
1 y	0.20 – 1.60																												
2 y	0.30 – 2.00																												
3 – 5 y	0.35 – 2.40																												
6 – 7 y	0.40 – 2.80																												
8 – 10 y	0.45– 3.20																												
11 – 15 y	0.50 – 3.80																												
16 – 150 y	0.60 – 4.20																												
<5 m	0.14 – 1.40																												
5 m- 3 y	0.20 – 1.60																												
4 –6 y	0.20 – 2.10																												
3 – 5 y	0.35 – 2.10																												
7 – 11 y	0.30 – 2.10																												
12 – 19 y	0.30 – 2.40																												
20 – 150 y	0.40 – 3.00																												
IRON, TOTAL IRON BINDING CAPACITY (TIBC), AND IRON SATURATION INDEX, S	<p>IRON $\mu\text{mol/L}$ MALE FEMALE U/X</p> <table> <tr><td>18 – 150 y</td><td>8 – 35</td></tr> </table> <p>TIBC $\mu\text{mol/L}$ MALE AND FEMALE</p> <table> <tr><td>0 – <18 y</td><td>50 – 80</td></tr> <tr><td>18 – 150 y</td><td>40 – 75</td></tr> </table> <p>IRON SATURATION INDEX: MALE FEMALE U/X</p> <table> <tr><td><18 y</td><td>0.10-0.50</td></tr> <tr><td>18 – 150 y M</td><td>0.12 – 0.60</td></tr> <tr><td>18-150 y F</td><td>0.10 -0.55</td></tr> <tr><td>18-150 U/X</td><td>0.10-0.60</td></tr> </table>	18 – 150 y	8 – 35	0 – <18 y	50 – 80	18 – 150 y	40 – 75	<18 y	0.10-0.50	18 – 150 y M	0.12 – 0.60	18-150 y F	0.10 -0.55	18-150 U/X	0.10-0.60														
18 – 150 y	8 – 35																												
0 – <18 y	50 – 80																												
18 – 150 y	40 – 75																												
<18 y	0.10-0.50																												
18 – 150 y M	0.12 – 0.60																												
18-150 y F	0.10 -0.55																												
18-150 U/X	0.10-0.60																												
LACTATE, Arterial	0.5 – 2.2 mmol/L																												
LACTATE, P	<p>MALE AND FEMALE mmol/L</p> <table> <tr><td>0 – 150 y</td><td>0.5 – 2.2</td></tr> </table>	0 – 150 y	0.5 – 2.2																										
0 – 150 y	0.5 – 2.2																												
LACTATE, CSF	<p>MALE AND FEMALE mmol/L</p> <table> <tr><td><3 d</td><td>1.1-6.7</td></tr> <tr><td>3 to < 10d</td><td>1.1-4.4</td></tr> <tr><td>10d to 150 y</td><td>1.1-2.4</td></tr> </table>	<3 d	1.1-6.7	3 to < 10d	1.1-4.4	10d to 150 y	1.1-2.4																						
<3 d	1.1-6.7																												
3 to < 10d	1.1-4.4																												
10d to 150 y	1.1-2.4																												
LACTOSE, P/S - TOLERANCE TEST - 2 h	<p>Lactose Tolerance Interpretive Criteria: Normal: Rise in glucose of >1.1 or greater mmol/L at any time post-lactose. Abnormal (query lactase deficiency): Rise in glucose of < =1.1 mmol/L at any time post-lactose.</p>																												
LACTATE DEHYDROGENASE (LD), P/S	<p>MALE AND FEMALE U/L</p> <table> <tr><td><1y</td><td>200-420</td></tr> <tr><td>1 to 9 y</td><td>140-320</td></tr> <tr><td>10 to 14 y</td><td>120-300</td></tr> <tr><td>15 to 150 y</td><td>120-250</td></tr> </table>	<1y	200-420	1 to 9 y	140-320	10 to 14 y	120-300	15 to 150 y	120-250																				
<1y	200-420																												
1 to 9 y	140-320																												
10 to 14 y	120-300																												
15 to 150 y	120-250																												

Test	Reference Intervals																				
LACTATE DEHYDROGENASE (LD), CSF	No reference range																				
LUTEINIZING HORMONE (LH), P/S	IU/L <table border="1" data-bbox="797 310 1531 785"> <thead> <tr> <th colspan="2" data-bbox="797 310 1531 346">Female and X/U</th> </tr> <tr> <th data-bbox="797 346 1036 382">Age</th> <th data-bbox="1036 346 1531 382">Interval</th> </tr> </thead> <tbody> <tr> <td data-bbox="797 382 1036 417"><11y</td> <td data-bbox="1036 382 1531 417"><7.0</td> </tr> <tr> <td data-bbox="797 417 1036 604">11-150 y</td> <td data-bbox="1036 417 1531 604"> See comment Follicular: 1.0 – 13.0 IU/L Luteal: 1.0 – 17.0 IU/L Midcycle: 8.0 – 76.0 IU/L Post-menopausal: 16.0-54.0 IU/L </td> </tr> <tr> <th colspan="2" data-bbox="797 604 1531 640">Male</th> </tr> <tr> <th data-bbox="797 640 1036 676">Age</th> <th data-bbox="1036 640 1531 676">Interval</th> </tr> <tr> <td data-bbox="797 676 1036 711"><11y</td> <td data-bbox="1036 676 1531 711"><7.0</td> </tr> <tr> <td data-bbox="797 711 1036 747">11y to <70y</td> <td data-bbox="1036 711 1531 747">1.0-9.0</td> </tr> <tr> <td data-bbox="797 747 1036 785">70y to 150 y</td> <td data-bbox="1036 747 1531 785">3.0-35.0</td> </tr> </tbody> </table>	Female and X/U		Age	Interval	<11y	<7.0	11-150 y	See comment Follicular: 1.0 – 13.0 IU/L Luteal: 1.0 – 17.0 IU/L Midcycle: 8.0 – 76.0 IU/L Post-menopausal: 16.0-54.0 IU/L	Male		Age	Interval	<11y	<7.0	11y to <70y	1.0-9.0	70y to 150 y	3.0-35.0		
Female and X/U																					
Age	Interval																				
<11y	<7.0																				
11-150 y	See comment Follicular: 1.0 – 13.0 IU/L Luteal: 1.0 – 17.0 IU/L Midcycle: 8.0 – 76.0 IU/L Post-menopausal: 16.0-54.0 IU/L																				
Male																					
Age	Interval																				
<11y	<7.0																				
11y to <70y	1.0-9.0																				
70y to 150 y	3.0-35.0																				
LIPASE, P/S	MALE AND FEMALE U/L 0-<18 y <50 18-150 y <80																				
LIPID PROFILE, P/S	<p data-bbox="773 909 1190 942"><u>Pediatric Lipid Profile (0 – 17 yr)</u></p> <table data-bbox="773 942 1292 1110"> <tr><td>Total Cholesterol</td><td>No reference range</td></tr> <tr><td>HDL-C</td><td>No reference range</td></tr> <tr><td>LDL-C</td><td>No reference range</td></tr> <tr><td>Triglycerides</td><td>No reference range</td></tr> <tr><td>Non-HDL-C</td><td>No reference range</td></tr> </table> <p data-bbox="773 1142 1531 1205">The following comment will be attached to the Non-HDL-C result:</p> <p data-bbox="773 1245 1515 1308">Lipid Profile acceptable limits relative to dyslipidemia and atherosclerosis risk:</p> <p data-bbox="773 1308 1336 1341">Total Cholesterol Acceptable <4.40 mmol/L</p> <p data-bbox="773 1341 1203 1375">HDL-C Acceptable >1.16 mmol/L</p> <p data-bbox="773 1375 1198 1409">LDL-C Acceptable <2.84 mmol/L</p> <p data-bbox="773 1409 1484 1472">Triglycerides Acceptable <0.85 mmol/L (0-9 years) OR <1.02 mmol/L (10-17 years)</p> <p data-bbox="773 1472 1162 1505">Non-HDL-C Acceptable <3.10</p> <p data-bbox="773 1545 1175 1579"><u>Adult Lipid Profile (18 – 150 yr)</u></p> <table data-bbox="773 1579 1292 1747"> <tr><td>Total Cholesterol</td><td>No reference range</td></tr> <tr><td>HDL-C</td><td>No reference range</td></tr> <tr><td>LDL-C</td><td>0.0 – 3.4 mmol/L</td></tr> <tr><td>Triglycerides</td><td>0.0 – 1.7 mmol/L</td></tr> <tr><td>Non-HDL-C</td><td>0.0 – 4.2 mmol/L</td></tr> </table> <p data-bbox="773 1778 1531 1841">The following comment will be attached to the Non-HDL-C result:</p> <p data-bbox="784 1881 1531 1980">For patients 30 years of age or older, the Framingham Risk Score (FRS), modified for family history, is recommended for risk assessment (2016 CCS Guideline,</p>	Total Cholesterol	No reference range	HDL-C	No reference range	LDL-C	No reference range	Triglycerides	No reference range	Non-HDL-C	No reference range	Total Cholesterol	No reference range	HDL-C	No reference range	LDL-C	0.0 – 3.4 mmol/L	Triglycerides	0.0 – 1.7 mmol/L	Non-HDL-C	0.0 – 4.2 mmol/L
Total Cholesterol	No reference range																				
HDL-C	No reference range																				
LDL-C	No reference range																				
Triglycerides	No reference range																				
Non-HDL-C	No reference range																				
Total Cholesterol	No reference range																				
HDL-C	No reference range																				
LDL-C	0.0 – 3.4 mmol/L																				
Triglycerides	0.0 – 1.7 mmol/L																				
Non-HDL-C	0.0 – 4.2 mmol/L																				

Test	Reference Intervals
	<p>Can J Cardiol 2016):</p> <p>FRS Calculation Resources can be found at https://myhealth.alberta.ca/Alberta/Pages/Heart-Disease-Risk-Calculator.aspx.</p> <p>Low Risk (FRS < 10%) Treatment advised if LDL-C \geq 5.0 mmol/L Treatment target: > 50% reduction LDL-C</p> <p>Intermediate Risk (FRS 10 - 19%) Treatment advised if LDL-C \geq 3.5 mmol/L OR Non-HDL-C \geq 4.3 mmol/L OR ApoB \geq 1.2 g/L;</p> <p>Consider treatment for men \geq 50 and women \geq 60 yrs with one additional CV risk factor</p> <p>Treatment targets: LDL-C < 2.0 mmol/L OR decrease by > 50% OR Non-HDL-C < 2.6 mmol/L OR ApoB < 0.8 g/L</p> <p>High Risk (FRS \geq 20% or presence of high risk features) Treatment advised in all patients</p> <p>Treatment targets: LDL-C < 2.0 mmol/L OR decrease by > 50% OR Non-HDL-C < 2.6 mmol/L OR ApoB < 0.8 g/L</p>
LITHIUM, S	<p>MALE AND FEMALE mmol/L</p> <p>0 – 150 y</p> <p>Suggested Range:</p> <p>Acute Mania Therapy 1.00 – 1.50</p> <p>Bipolar Maintenance Therapy 0.60 – 1.20</p>
MAGNESIUM, 24 Hour Urine	<p>MALE AND FEMALE mmol/d</p> <p>0 – 150 y 2.00 -8.00</p>
MAGNESIUM, P/S	<p>MALE, FEMALE, U/Xmmol/L</p> <p>0-150y 0.70-1.00</p>
MALARIAL PARASITE SCREEN, B	Negative
MONONUCLEOSIS TEST, S	Negative
OCCULT BLOOD, F	Negative
OSMOLALITY, P/S	<p>MALE AND FEMALE mmol/kg</p> <p>0 – 150 y 280 – 300</p>
OSMOLALITY, U	<p>MALE AND FEMALE mmol/kg</p> <p>0 – 150 Y 50 – 1400</p>
pH, Arterial – Cord Blood	pH 7.20 - 7.40
pH, Venous – Cord Blood	pH 7.30 - 7.40
PHENOBARBITAL, P/S	<p>MALE AND FEMALE μmol/L</p> <p>Therapeutic Range</p> <p>0 – 150 y 45 – 170</p>
PHENYTOIN, P/S	<p>MALE AND FEMALE μmol/L</p> <p>Therapeutic Range</p> <p><3 m 25 - 55</p> <p>3 m – 150 y 40 – 80</p>

Test	Reference Intervals
PHOSPHATE, 24 Hour Urine	MALE AND FEMALE mmol/d 0 – 150 yr 15.0-50.0
PHOSPHATE, P/S	MALE, FEMALE and U/X mmol/L 0 – 15 d 1.40 – 2.70 15 d – <1 m 1.60 – 2.70 1 m – <5 y 1.20 – 2.20 5 – <13 y 1.10 – 1.90 13 – <18 y 0.90 – 1.70 18 – 150 y 0.70 – 1.50
PROLACTIN, P/S	MALE ug/L 0 – 150 yr 4.0-15.0 FEMALE ug/L 0 – 150 y 4.0-25.0 X/U 4.0-25.0
PROTEIN – TOTAL, 24 Hour Urine	MALE AND FEMALE g/d 0 – 150 y < 0.20
PROTEIN – TOTAL, CSF	MALE AND FEMALE g/L 0 – <31d 0.14 – 1.12 31 d– 150 y 0.15 – 0.45
PROTEIN – TOTAL, P/S	MALE, FEMALE and UNKNOWN g/L 0 – 364 d 40 – 70 365 d – 150 y 62 – 82
PROTEIN – TOTAL, U	MALE AND FEMALE g/L 0 – 150 yr < 0.20
PROTEIN CREATININE RATIO, U	MALE AND FEMALE mg/mmol 2- 12 y 2.0-20.0 13-150 y <13.0
PT (PROTHROMBIN TIME), INR, P	0.8 - 1.2 Therapeutic Range: 2.0 - 3.0 NOTE: For some indications a higher target therapeutic range is required. Reference: February 2012, 141(2_suppl) Antithrombotic Therapy and Prevention of Thrombosis, 9 th ed: American College of Chest Physicians Evidence-Based Practice Guidelines.
PTT (PARTIAL THROMBOPLASTIN TIME), P	Reference Range and Therapeutic Range varies with instrument. Please refer to patient report for current value.
RETICULOCYTE COUNT, B	RELATIVE (%): 0 – 1 mo 2.0 – 6.0 1 mo – 11 y 0.9 – 2.7 12y – 150 yr (F) 1.2 – 2.7 12y – 150 yr(M) 0.9 – 2.1 ABSOLUTE: Birth 78 – 110 x 10 ⁹ /L 1d – 1wk 78 – 396 x 10 ⁹ /L 1wk – 1m 60 – 372 x 10 ⁹ /L 1m – 6m 24 – 132 x 10 ⁹ /L 6m – 5y 33 – 143 x 10 ⁹ /L

Test	Reference Intervals
	6y – 11y 36 – 140 x 10 ⁹ /L 12y – 17y (F) 47 – 138 x 10 ⁹ /L 12y – 17y (M) 39 – 111 x 10 ⁹ /L 18 – 150 yr (F) 49 – 140 x 10 ⁹ /L 18 – 150 yr (M) 42 – 122 x 10 ⁹ /L
RHEUMATOID FACTOR QUANTITATIVE, P/S	MALE AND FEMALE KU/L 0 – 150 yr < 21
ROTAVIRUS	Negative
RSV (RESPIRATORY SYNCYTIAL VIRUS)	Negative
RSV and INFLUENZA A/B	Negative
SALICYLATE, P/S	No reference range
SEMINAL FLUID ANALYSIS, FERTILITY, Semf	Liquefaction Normal (complete within 60 minutes of collection) Volume > 1.4 mL Viscosity Normal (< 2 cm thread) pH > 7.1 Sperm concentration > 14.9 M/mL Total sperm count > 38.9 M RBC 0/hpf WBC 0/hpf Sperm Motility > 31% Progressive or > 39% Progressive and Non-progressive within 60 minutes of ejaculation Vitality > 57% live
SEMINAL FLUID ANALYSIS, POST VASECTOMY, Semf	None seen
T3 – FREE, P/S	MALE AND FEMALE pmol/L 0 – 29 d 4.2-13.0 30 d – <1 y 5.1-8.6 1 – 13 y 4.4-8.1 14 – 17 y 3.5-7.4 18 – 150 y 3.0– 6.5
THROMBIN TIME, P	Reference interval varies with instrument/testing location. Please refer to the patient report or look in NetCare or the EMR.
THYROXINE FREE (T4), P/S	MALE AND FEMALE pmol/L 0 – 14 d 13.5 – 50.0 15 d – 29d 8.7-32.5 ≥30d 10.0-25.0
TOBRAMYCIN P/S	mg/L Pre-dose: <2.0 Post-dose: 5.0-10.0 8 hr interval: comment
TRIGLYCERIDE, P/S	MALE AND FEMALE: mmol/L 0 – 17 y < 1.50 The following comment will be attached to the result: Acceptable limit relative to dyslipidemia and atherosclerosis risk <0.85 mmol/L (0-9 years) OR <1.02

Test	Reference Intervals
	mmol/L (10-17 years). 18 – 150 y 0.00 – 1.70
TROPONIN I (Tnl), P/S	Reference range varies with instrument. Please refer to patient report for current value(s).
HS TROPONIN T P/S	MALE FEMALE <14 ng/L
TSH (THYROID STIMULATING HORMONE), P/S	MALE AND FEMALE mU/L 0 – 7d 1.23 – 25.00 30 d – <1y 1.00 – 6.80 >=1 y 0.20-6.50
TSH PROGRESSIVE (THYROID STIMULATING HORMONE), P/S	MALE AND FEMALE mU/L 0 – 29 d 1.23 – 25.00 8 d – <1y 1.00 – 6.80 >=1 y 0.20-6.50
URATE, 24 Hour Urine	MALE AND FEMALE mmol/d 0 – 150 y 1.5 – 4.5
URATE, P/S	MALE FEMALE U/X µmol/L <5y 100 – 300 5 y to <10 y 140 – 330 10– <18 y M 160-500 18-150y M 200-500 10 – <70y F 150-400 70-150y F 150-500 10-150y U/X 150-500
UREA, 24 Hour Urine	MALE AND FEMALE mmol/d 0 – 150 y 430-710
UREA, P/S	MALE, FEMALE, U/Xmmol/L <2 y 1.0 – 7.5 2 – <18y 2.0 – 7.0 MALE 18-<55 y 3.0 – 8.0 >55 y 3.0 – 9.0 FEMALE 18-<55 y 2.0– 7.0 >55 y 3.0 – 8.0 U/X 18-<55 2.0-8.0 >55 3.0-9.0
URINALYSIS, U	MALE & FEMALE 0 – 150 y Blood: Negative Clarity: Clear Color: Colorless, Yellow, Amber

Test	Reference Intervals
	Glucose: Negative Ketone: Negative Leukocytes: Negative Nitrite: Negative pH: 5 – 8 Protein: Negative SG: 1.005 – 1.030
URINALYSIS MICROSCOPIC, U	MALE AND FEMALE 0 – 150 yr WBC: 0 – 5/hpf RBC: 0 – 2/hpf Squamous/Transitional epithelial cells: 0 – 5/hpf Renal epithelial cells: Negative Bacteria: 0-20/ hpf Casts, hyaline: 0 – 2/lpf All other cast types: Negative Oval Fat Bodies, Trichomonas and Yeast: Negative
VALPROATE (VALPROIC ACID), P/S	MALE AND FEMALE $\mu\text{mol/L}$ 0 – 150 y Therapeutic Range 350 – 700 NOTE: Concentrations up to 1040 $\mu\text{mol/L}$ may be required for some patients with complex partial seizures and secondarily generalized tonic-clonic seizures.
VANCOMYCIN, P - RANDOM	No range
VANCOMYCIN, P – TROUGH PRE	MALE AND FEMALE mg/L Therapeutic Range 0 – 150 y 10.0 – 20.0
VITAMIN B12, P/S	MALE, FEMALE & U/X pmol/L 0 – <10 y ≥ 250 10-150 y ≥ 160

Blood Gases and Co-oximetry Reference Intervals

Test Name	Adult / Pediatric			
	Arterial	Venous	Mixed Venous	Capillary
pH	7.35 – 7.45	7.32 - 7.42	7.33 – 7.43	7.34 - 7.44
pCO ₂ (mmHg)	35 – 45	40 - 50	37 – 47	37 – 47
pO ₂ (mmHg)	70 – 90	30 - 50	35 – 45	40 – 60
HCO ₃ (mmol/L)	20 – 26	22 - 28	21 – 27	21 – 27
tCO ₂ (mmol/L)	21 – 28	23 - 30	22 – 29	22 – 29
Base Excess (mmol/L)	- 2 to +2	- 2 to +2	-2 to +2	- 2 to +2
O ₂ Saturation (%)	92 – 98	60 - 80	70 – 80	90 – 95
tHb (g/L)	120 – 180	120 – 180		120 – 180
O ₂ Hb (%)	92 – 98	64 – 74		90 – 95
COHb (%)	0 – 3	0 – 3		0 – 3
MetHb (%)	0 – 1.5	0 – 1.5		0 – 1.5
HHb (%)	0 – 5	23 – 33		0 – 7
O ₂ CT (mL/dL)	16 – 24	7 – 18		13 – 22
Aa DO ₂ (mmHg)	<15 Room Air <100 100% O ₂			

Test Name	Cord Blood	
	Arterial Cord Blood	Venous Cord Blood
pH	7.20 – 7.40	7.25 - 7.45
pCO ₂ (mmHg)	32 – 66	27 – 49
pO ₂ (mmHg)	6 – 30	17 – 41
HCO ₃ (mmol/L)	17 – 27	16 – 25
tCO ₂ (mmol/L)	19 – 29	18 – 27
Base Excess (mmol/L)	- 8 to +2	- 8 to +2
O ₂ Saturation (%)	5 – 60	15 – 75