

Form Title **Diabetic Ketoacidosis Pediatric Inpatient Order Set
(for Sites Using D10W Solutions)**

Form Number **21165Bond**

© 2018, Alberta Health Services, CKCM



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. The license does not apply to content for which the Alberta Health Services is not the copyright owner.

To view a copy of this license, visit

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Disclaimer: This material is intended for use by clinicians only and is provided on an "as is", "where is" basis. Although reasonable efforts were made to confirm the accuracy of the information, Alberta Health Services does not make any representation or warranty, express, implied or statutory, as to the accuracy, reliability, completeness, applicability or fitness for a particular purpose of such information. This material is not a substitute for the advice of a qualified health professional. Alberta Health Services expressly disclaims all liability for the use of these materials, and for any claims, actions, demands or suits arising from such use.

**Diabetic Ketoacidosis Pediatric Inpatient
Order Set (for Sites Using D10W Solutions)**

Refer to Diabetic Ketoacidosis, Pediatric - Emergency and Inpatient for the Pediatric DKA Management Algorithm
Select orders by placing a (✓) in the associated box
Physician _____

Last Name (Legal)		First Name (Legal)	
Preferred Name <input type="checkbox"/> Last <input type="checkbox"/> First		DOB(dd-Mon-yyyy)	
PHN	ULI <input type="checkbox"/> Same as PHN	MRN	
Administrative Gender <input type="checkbox"/> Male		<input type="checkbox"/> Female	
<input type="checkbox"/> Non-binary/Prefer not to disclose (X)		<input type="checkbox"/> Unknown	

Weight _____ Kg

Admission/ Discharge/ Transfer

Admit to inpatient unit under care of Dr. _____

Patient Care

Weigh patient, daily

Diet

NPO

Sugar free oral fluids when serum bicarbonate greater than or equal to 18 mmol/L
(mild acidosis/ketosis may still be present).

Monitoring

Vital signs and neurovitals are indicated at a minimum of every hour in the initial 1-4 hours; more frequently as indicated (based on severity and care setting)

Notify physician if:

- Decreased or changing level of consciousness (*restless, irritable, drowsy, obtunded*) especially after initial improvement
- Headache, hypertension, vomiting, incontinence, cranial nerve palsies, oxygen desaturation

Initial (for 4 hours)

- Vital signs: heart rate, blood pressure, respiratory rate, temperature; every _____ hour
- Neurovitals: level of consciousness, Glasgow coma scale (GCS) every _____ hour
- Vital signs: Continuous Pulse oximetry or cardiac monitor
- Intake and Output: Strictly monitor intake and output hourly

After clinical improvement

Vital signs and neurovitals are indicated every 2 to 4 hours if DKA is improving and level of consciousness is normal

- Vital signs every _____ hours
- Neuro vital signs every _____ hours
- Intake and Output: Monitor fluid volume intake and output every 4 hours

Point of Care Testing

Frequent blood glucose point of care testing is required while adjusting insulin/IV in first 1-4 hours of presentation; may be reduced to every 2 hours if blood glucose is stable

Initial (until blood glucose stable at 10 to 15 mmol/L for 4 hours)

- Blood Glucose Monitoring - POCT, by finger poke every hour AND PRN; *Check blood glucose using glucometer at the bedside prior to administering any IV fluids*
- Urine Ketones - POCT with each void (*optional: if testing readily available*)

After clinical improvement (once blood glucose has been stable at 10 to 15 mmol/L for 4 hours)

- Blood Glucose Monitoring - POCT, by finger poke; every _____ hours AND PRN by Glucometer
- Discontinue Urine Ketones POCT when ketones negative for _____ void

Prescriber Name	Prescriber Signature	Date (dd-Mon-yyyy)	Time (hh:mm)
-----------------	----------------------	--------------------	--------------

**Diabetic Ketoacidosis Pediatric Inpatient
Order Set (for Sites Using D10W Solutions)**

Physician _____

Last Name (Legal)		First Name (Legal)	
Preferred Name <input type="checkbox"/> Last <input type="checkbox"/> First		DOB (dd-Mon-yyyy)	
PHN	ULI <input type="checkbox"/> Same as PHN	MRN	
Administrative Gender <input type="checkbox"/> Male		<input type="checkbox"/> Female	
<input type="checkbox"/> Non-binary/Prefer not to disclose (X)		<input type="checkbox"/> Unknown	

Laboratory Investigations Repeating
Chemistry

- Hemoglobin A1C (if not done in Emergency or within the last 30 days)
- Obtain and repeat the following labs every _____ hours
(The following are recommended at a minimum of every 4 hours to monitor response to therapy)

- Sodium (Na)
- Potassium (K)
- Chloride (Cl)
- Glucose Random
- Bicarbonate

- Repeat the following labs every _____ hours
(the following are recommended at a minimum of every 8 hour to monitor response to therapy)

- Serum osmolality
- Creatinine
- Urea
- Calcium
- Beta-hydroxybutyrate
- Phosphate
- Magnesium
- Discontinue lab investigations when insulin infusion has been discontinued.

Blood Gases

Capillary or venous blood gases are acceptable.

- Blood gas capillary every 4 hours
 (optional) Alternate collection with chemistry every 4 hours to monitor labs every 2 hours
- Blood gas venous every 4 hours
 (optional) Alternate collection with chemistry every 4 hours to monitor labs every 2 hours
- Ionized calcium every 4 hours (if not included with blood gas result)

Fluid Management

Proceed to detailed re-hydration calculations after the first hour of initial fluid management (see attached) Proceed to detailed rehydration calculations. Until detailed fluid calculations are completed continue initial IV infusion rate (1.5 x maintenance for mild/ moderate DKA and 2 x maintenance for severe DKA).

Avoid over hydration, total fluid should not exceed 2X maintenance

To avoid inadequate rehydration and in the context of challenging hydration assessment in DKA:

- Used attached calculation sheet based on mild, moderate or severe DKA
- Order total hourly fluid rate of Bag A and Bag B based on this calculation

Prescriber Name	Prescriber Signature	Date (dd-Mon-yyyy)	Time (hh:mm)
-----------------	----------------------	--------------------	--------------

**Diabetic Ketoacidosis Pediatric Inpatient
Order Set (for Sites Using D10W Solutions)**

Last Name (Legal)		First Name (Legal)	
Preferred Name <input type="checkbox"/> Last <input type="checkbox"/> First		DOB(dd-Mon-yyyy)	
PHN	ULI <input type="checkbox"/> Same as PHN	MRN	
Administrative Gender <input type="checkbox"/> Male <input type="checkbox"/> Female		<input type="checkbox"/> Non-binary/Prefer not to disclose (X) <input type="checkbox"/> Unknown	

Physician _____

Fluid Management (Continued)						
<p>Total Hourly Fluid Rate based on Detailed Fluid Calculations: See Attached</p> <p><input checked="" type="checkbox"/> Total IV fluids (excluding insulin) = _____ mL/hr for the next 48 hours or until IV is discontinued</p> <p>• Total Hourly Fluid Rate = Bag A (saline) + Bag B (saline and dextrose)</p> <p><input checked="" type="checkbox"/> IV should be reduced by the amount of oral intake once oral fluids are ordered to maintain total hourly fluid rate.</p>						
<p>1. Initiate Bag A (Use when blood glucose greater than 17 mmol/L. Once blood glucose approaches 17 move to Bag A + Bag B system)</p> <p><i>If additional potassium is required in addition to 40 mEq/L potassium in IV fluids, follow local policy regarding availability of IV fluids containing 60 mEq/L of KCL</i></p> <p><input type="checkbox"/> 0.9% NaCl with _____ mEq/L KCL (recommended 40 mEq/L) at calculated total hourly fluid rate = _____ mL/hour</p>						
<p>2. Add Bag B once blood glucose is less than or equal to 17 mmol/L</p> <p><input checked="" type="checkbox"/> 10% Dextrose System with 0.9% NaCl</p> <p><i>Start with a combination of Bag A and Bag B that provides a dextrose concentration of D10W/0.9% NaCl with 40 mEq KCL</i></p> <p>This is accomplished by:</p> <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:50%; vertical-align: top;"> <ul style="list-style-type: none"> • Bag A: 0.9% NaCl with _____ mEq/L KCL <i>(recommended 40 mEq/L)</i> • Rate: (0% of total hourly fluid rate = total hourly fluid rate X 0) = _____ mL/hour </td> <td style="width:10%; text-align: center; vertical-align: middle;">AND</td> <td style="width:50%; vertical-align: top;"> <ul style="list-style-type: none"> • Bag B: D10W/0.9% NaCl with _____ mEq/L KCL <i>(recommended 40 mEq/L)</i> • Rate: (100% of total hourly fluid rate = total hourly fluid rate X 1) = _____ mL/hour </td> </tr> </table>				<ul style="list-style-type: none"> • Bag A: 0.9% NaCl with _____ mEq/L KCL <i>(recommended 40 mEq/L)</i> • Rate: (0% of total hourly fluid rate = total hourly fluid rate X 0) = _____ mL/hour 	AND	<ul style="list-style-type: none"> • Bag B: D10W/0.9% NaCl with _____ mEq/L KCL <i>(recommended 40 mEq/L)</i> • Rate: (100% of total hourly fluid rate = total hourly fluid rate X 1) = _____ mL/hour
<ul style="list-style-type: none"> • Bag A: 0.9% NaCl with _____ mEq/L KCL <i>(recommended 40 mEq/L)</i> • Rate: (0% of total hourly fluid rate = total hourly fluid rate X 0) = _____ mL/hour 	AND	<ul style="list-style-type: none"> • Bag B: D10W/0.9% NaCl with _____ mEq/L KCL <i>(recommended 40 mEq/L)</i> • Rate: (100% of total hourly fluid rate = total hourly fluid rate X 1) = _____ mL/hour 				
<p><input checked="" type="checkbox"/> Titrate dextrose infusion to maintain blood glucose 8-15mmol/L. Adjust with each hourly blood glucose level (by Blood Glucose Monitoring POCT)</p> <p>If blood glucose is greater than 15 mmol/L → Increase Bag A (saline) rate and decrease Bag B (saline and dextrose) rate by 25% of total hourly fluid rate</p> <p>If blood glucose is less than 10 mmol/L → Decrease Bag A rate (saline) and increase Bag B rate (saline and dextrose) by 25% of total hourly fluid rate</p> <p>If blood glucose is between 10 to 15 mmol/L → No changes to IV rates for either solution.</p> <p>Total hourly fluid rate _____ mL/hour X 0.20 = _____ mL/hour (Amount by which the the IV fluid rate of each bag will change with each adjustment, maintaining total hourly fluid rate)</p> <p><input type="checkbox"/> If blood glucose decreases more than 5 mmol/L per hour, contact physician.</p>						
Prescriber Name	Prescriber Signature	Date (dd-Mon-yyyy)	Time (hh:mm)			

**Diabetic Ketoacidosis Pediatric Inpatient
Order Set (for Sites Using D10W Solutions)**

Physician _____

Last Name (Legal)		First Name (Legal)	
Preferred Name <input type="checkbox"/> Last <input type="checkbox"/> First		DOB (dd-Mon-yyyy)	
PHN	ULI <input type="checkbox"/> Same as PHN	MRN	
Administrative Gender <input type="checkbox"/> Male		<input type="checkbox"/> Female	
<input type="checkbox"/> Non-binary/Prefer not to disclose (X)		<input type="checkbox"/> Unknown	

Medications
Insulin Infusion

To be ordered after patient has received 1 to 2 hours of IV fluids and is hemodynamically stable.

If metabolic acidosis is not improving after 4 hours, re-evaluate to ensure rehydration calculations are correct, insulin infusion is properly mixed, and that intravenous are not occluded, are patent and infusing. Once these are re-evaluated, if no improvement consider consulting pediatric endocrinology and/or PICU.

Humulin Regular insulin (1 unit/mL in 0.9% NaCl) at a starting dose of 0.1 units/kg/hour = _____ units/hour
= _____ mL/hour intravenous

Additional DKA orders as required
Oral Potassium Supplementation

If additional potassium is required in addition to 40 Eq/L potassium in IV fluids, follow local policy if available regarding availability of IV fluids containing 60 mEq/L of KCL, or consider oral potassium supplementation (may cause vomiting).

Potassium Chloride _____ mEq (1 mEq/kg/dose) PO every 12 hours for _____ doses (if patient has normal level of consciousness, even if NPO. 1- 3 doses recommended)

Ongoing Fluid Management

For most patients 0.9% NaCl can be used with DKA resolution. If a decrease to 0.45% NaCl is required (e.g. in the case of hyperchloremia) it should only be considered if corrected sodium is 140 to 150 mmol/L and stable and patient has received 4 to 5 hours of fluid.

If additional potassium is required in addition to 40 Eq/L potassium in IV fluids, follow local policy regarding availability of IV fluids containing 60 mEq/L of KCL, or consider oral potassium supplementation (may cause vomiting).

- Change Bag A to 0.45% NaCl with _____ mEq KCl/L
- Change Bag B to D10W/0.45% NaCl with _____ mEq KCl/L
- Continue titration of Bag A and Bag B as established on page 3

For infusion rate change orders as required, contact the most responsible practitioner

Phosphate

If serum phosphate is less than 0.4 mmol/L, consider administering in a separate sodium phosphate (NaPO₄) infusion if possible. If pre-mixed NaPO₄ IV solution bags are not available, an IV infusion of NaPO₄ can be prepared by using vials of sodium phosphate. Mixing and administration instructions are on Insite. Use local infusion protocols and order sets if available.

If phosphate is given, monitor serum Ca, Mg, and PO₄ levels minimum every 4 hours to avoid hypocalcemia.

Prescriber Name	Prescriber Signature	Date (dd-Mon-yyyy)	Time (hh:mm)

**Diabetic Ketoacidosis Pediatric Inpatient
Order Set (for Sites Using D10W Solutions)**

Last Name (Legal)		First Name (Legal)	
Preferred Name <input type="checkbox"/> Last <input type="checkbox"/> First		DOB (dd-Mon-yyyy)	
PHN	ULI <input type="checkbox"/> Same as PHN	MRN	
Administrative Gender <input type="checkbox"/> Male <input type="checkbox"/> Female		<input type="checkbox"/> Non-binary/Prefer not to disclose (X) <input type="checkbox"/> Unknown	

Physician _____

Detailed Fluid Calculations				
Body Weight in kilograms:		kg (a)		
Step	Clinical Decision Support			Calculation
1. Identify percentage dehydration based on severity of DKA <i>Refer to Diabetic Ketoacidosis, Pediatric - Emergency and Inpatient for the severity of DKA</i>	Severity of DKA	% Dehydration	Extent of Dehydration (mL/kg)	Extent of dehydration = _____ mL/kg (b)
	Mild to Moderate DKA	6%	60mL/kg	
	Severe DKA	9%	90mL/kg	
2. Calculate total fluid deficit	Multiply: (body weight in kg) x (extent of dehydration in mL/kg)			Body weight (a) X dehydration mL/kg (b) = Total fluid deficit _____ mL (i)
3. Calculate the remainder of the fluid deficit (ii) after the initial 1-2 hours of fluid management	Amount of IV fluid received in first 1-2 hours: _____ mL (c) Subtract: [Total fluid deficit (i)] - [amount NS given first 1-2 hrs (c)]			(i) - (c) = _____ mL (ii) = Fluid deficit (ii) to be given over the next 48 hours in addition to maintenance fluids
4. Calculate maintenance fluid requirements for the next 48 hours	24-hour maintenance requirements based on body weight: 10 kg or less = 100 mL/kg/24 hours 11–20 kg = 1000 mL + 50 mL/kg/24 hours for each kg greater than 10 Greater than 20 kg = 1500 mL+20 mL/kg/24 hour for each kg greater than 20 = _____ mL maintenance for 24 hours Multiply by 2 = _____ 48 hour maintenance (iii)			24 hour maintenance requirements X 2 = _____ mL (iii) = Maintenance fluid requirements for the next 48 hours
5. Calculate the total amount of fluid still to be given over 48 hours	Ongoing fluid losses (i.e gastric losses): _____ mL (iv) Total Fluid to be given over 48 hours Add: FLUID DEFICIT (ii) + MAINTENANCE FLUIDS (iii) <i>Note: Additional Losses (ie vomiting/ diarrhea) are NOT included; If added losses are occurring, measure losses and replace in addition.</i>			(ii) + (iii) = _____ mL (v) = Total Amount of Fluid to be given in next 48 hours (v)
6. Calculate hourly fluid rate for the next 48 hours.	Total amount of fluid to be given in 48 hours mL (v) /48 hours = hourly rate of fluid replacement (in mLs/hr) **Note: If calculated fluid deficit replacement rate PLUS maintenance is greater than 2x maintenance ie in severe DKA in a larger child) use a maximum rate of 2x maintenance.			(iv) mL/48 hours = _____ mL/hr = Total Hourly Fluid Rate (**maximum rate is 2x 48 hour maintenance)