



For use at Alberta Health Services (AHS) immunization programs.

Use this form **when a parent or alternate decision-maker is not able** to be with the person being immunized at an AHS immunization service. *(An alternate decision-maker could be an agent, guardian, specific decision-maker or co-decision-maker.)*

The parent or alternate decision-maker should complete this form and send it - together with a copy of the documents showing they are authorized as alternate decision-maker - with the person being immunized, to deliver to the AHS immunization service. **These documents are required for this person to be immunized.**

Personal information <i>(of person being immunized)</i>	
Name <i>(Last, First, Middle)</i>	
Date of Birth <i>(dd-Mon-yyyy)</i>	Personal Health Number
Health information for the person being immunized <i>(If you need more space, use Notes section on Page 2, Side A)</i>	
Does this person have any allergies, including allergies to any vaccine, medicine, or food? <input type="checkbox"/> No <input type="checkbox"/> Yes ► List all allergies and reactions	
Does this person have any chronic <i>(long-term)</i> health conditions? <input type="checkbox"/> No <input type="checkbox"/> Yes ► List all health conditions	
If this person is under age 18 years, have they ever had multisystem inflammatory syndrome (MIS-C)? <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> Not applicable	
Is this person taking any medicine? <input type="checkbox"/> No <input type="checkbox"/> Yes ► List all medications	
Is this person pregnant? <input type="checkbox"/> No <input type="checkbox"/> Yes	Is this person breastfeeding/chestfeeding? <input type="checkbox"/> No <input type="checkbox"/> Yes
Has this person had COVID-19 vaccine before? <input type="checkbox"/> No <input type="checkbox"/> Yes ► List vaccine(s) and date(s)	
Has this person ever had a side effect from COVID-19 immunization? <input type="checkbox"/> No <input type="checkbox"/> Yes ► <i>(describe)</i> _____	
Has this person had myocarditis or pericarditis within 6 weeks of getting a COVID-19 vaccine? <input type="checkbox"/> No <input type="checkbox"/> Yes	
Has this person had a COVID-19 infection? <input type="checkbox"/> No <input type="checkbox"/> Yes ► When was their infection? _____	
Has this person had any vaccines in the last 2 weeks? <input type="checkbox"/> No <input type="checkbox"/> Yes ► List vaccine(s) and date(s)	

Alberta Health Services collects health information according to Section 20 of the Health Information Act (HIA). This information is used to provide health services, determine eligibility for health services, or to carry out any other purpose authorized by the HIA. If you have any questions about this, please ask the healthcare provider giving the immunization or contact your local public health and community health centre. If you do not know your local public health and community health centre, call Health Link at 811 to get this information.

Person Being Immunized

 Name *(Last, First, Middle)*

 Date of Birth *(dd-Mon-yyyy)*

Personal Health Number

Consent

I confirm that I have read the attached COVID-19 vaccine information.

- I know about and understand the risks, benefits, and common side effects of this vaccine.
- I know I can contact my local public health or community health centre or call Health Link at 811 if I have any questions. At the time of signing this form, I have no questions about this person getting this vaccine.
- I understand this consent is for all doses of the vaccine.
- I will contact my local public health or community health centre or the healthcare provider giving the COVID-19 vaccine before this person gets any dose if:
 - their health changes
 - they have a severe or unusual side effect after getting the COVID-19 vaccine *(other than the expected side effects listed on the COVID-19 vaccine information sheet)*
- I understand the information I have been given.
- I confirm that I have the legal authority to consent to this person's immunization.

 I consent to this person getting *(choose one only)*:

- Either** the Moderna Spikevax KP.2 **or** the Pfizer-BioNTech Comirnaty KP.2 mRNA COVID-19 vaccine
(Depending which vaccine is recommended for this person at the time of their immunization)
- Moderna Spikevax KP.2 mRNA COVID-19 vaccine *(for ages 6 months and older)*
- Pfizer-BioNTech Comirnaty KP.2 mRNA COVID-19 vaccine *(for ages 12 years and older)*

 I understand that I may withdraw *(take back)* this consent at any time by calling the local public health or community health centre or healthcare provider giving this vaccine.

 Name of Person Giving Consent *(print)*

Signature of Person Giving Consent

Relationship to person

- Parent (with legal authority to consent) Guardian/Legal representative
- Agent Co-decision-maker
- Specific decision-maker Other _____

 Date *(dd-Mon-yyyy)*

Daytime Phone

Other Phone

Person Being Immunized

Name (*Last, First, Middle*)

Date of Birth (*dd-Mon-yyyy*)

Personal Health Number

For Office Use Only

Phone/Fax Consent

Mode Consent was Received Fax/Scan Phone

Name of Healthcare Provider obtaining the consent

Date (*dd-Mon-yyyy*)

Time (*24hrs*)

Signature of Healthcare Provider obtaining the consent

Consent Using an Interpreter (*for non-English speaking parent/client*)

Interpreter's name or ID #

Phone

Date (*dd-Mon-yyyy*)

Time (*24hrs*)

Notes (*for Office Use, or if more space for health information is needed*)

This page is intentionally blank.

COVID-19 vaccines – mRNA

(Age 6 months and older)

What are COVID-19 vaccines?

COVID-19 vaccines protect against the SARS-CoV-2 virus (also known as COVID-19). The virus causes an infection in the lungs and airways and is a type of respiratory illness. In some cases, the infection can cause problems with other organs or other parts of the body. Go to ahs.ca/covid to learn more about COVID-19.

COVID-19 mRNA vaccines help protect you against getting very sick from COVID-19 and needing care in the hospital. The Moderna SpikeVax KP.2 and Pfizer-BioNTech Comirnaty KP.2 vaccines are available in Alberta. They are updated mRNA vaccines that help protect you against the COVID-19 strains that are currently spreading.

Who should get an mRNA COVID-19 vaccine?

Everyone is at risk of COVID-19. COVID-19 vaccines are free and recommended for everyone age 6 months and older.

Immunization is strongly recommended if you are at higher risk of getting COVID-19 or getting very sick from COVID-19. This includes if:

- You are age 65 years or older.
- You live in a continuing care or supportive living facility.
- You have a weak immune system.
- You are an Indigenous person.
- You are pregnant.
- You are a member of a racialized and other equity deserving community.
- You are a person who provides essential community services, including healthcare workers.

Do I need a vaccine if I have already had COVID-19?

If you have already had COVID-19, it is not yet known how long your protection will last or how much protection you will have against variants. It is important to get a COVID-19 vaccine even if you have already had the virus.

The vaccine may give you better protection if you wait a while after having COVID-19 and then get a vaccine. How long to wait depends on your health history, the number of doses of COVID-19 vaccine you have had, and your risk of getting very sick from COVID-19. If you have had COVID-19 in the past, check with your healthcare provider about when to get a COVID-19 vaccine.

What if my child is getting a COVID-19 vaccine?

A parent or guardian provides consent for children under age 18 years to get a COVID-19 vaccine. If a parent or guardian cannot be at the appointment, they can give consent in writing using the consent form at ahs.ca/VaccineUnder18. In some cases, children under age 18 years may be able to give their own consent.

How well do COVID-19 vaccines work?

How well COVID-19 vaccines work is different for each variant (also called strain) of the virus. As new COVID-19 variants are identified, vaccines are updated to offer better protection.

The mRNA vaccines give the best protection against COVID-19. They help to lower your risk of getting very sick from COVID-19 and needing care in the hospital.

What vaccine can I get?

If you are age 6 months to 11 years, you can get the Moderna vaccine.

If you are age 12 years or older, you can get either the Moderna vaccine or the Pfizer-BioNTech vaccine.

How many doses of mRNA COVID-19 vaccine do I need?

6 months to 4 years with a healthy immune system

If you are age 6 months to 4 years, you need 2 doses of a COVID-19 vaccine, 8 weeks apart. The Moderna KP.2 vaccine can start or complete your 2-dose series.

If you have already had 2 or more doses of a non-KP.2 COVID-19 vaccine, you can get 1 dose of the Moderna KP.2 vaccine at least 3 months after your last dose of COVID-19 vaccine.

5 years and older with a healthy immune system

If you are age 5 years or older, you need 1 dose of a KP.2 COVID-19 vaccine. If you have had another COVID-19 vaccine in the past, you need to wait at least 3 months before getting a KP.2 COVID-19 vaccine.

Weak immune system

If you have a weak immune system, the number of doses you need depends on your age and how many COVID-19 vaccines you have had in the past.

You may have a weak immune system if:

- You have had or will have an organ or stem cell transplant.
- You have had CAR T-cell therapy (a type of cancer treatment)
- You have kidney disease and need dialysis.
- You have human immunodeficiency virus (HIV) infection or acquired immunodeficiency syndrome (AIDS).
- You take certain medicines that weaken your immune system.

If you have a weak immune system:

- You need 3 doses of KP.2 COVID-19 vaccine if you have never had any COVID-19 vaccine.
- You need 1 dose of KP.2 vaccine at least 3 months after your last dose if you have already had 3 or more doses of a non-KP.2 vaccine.
- You need 1 or 2 doses of KP.2 vaccine to total a 3-dose series if you have already had 1 or 2 doses of a non-KP.2 vaccine.

Studies show that extra doses may give better protection to adolescents and adults with a weak immune system. In babies and children with a weak immune system, extra doses may also give better protection, but research is still happening to learn more. Research has shown mRNA vaccines provide the best protection for people with a weak immune system.

If you have had a stem cell transplant or are getting CAR T-cell therapy, talk to your healthcare provider about when you can get a dose of KP.2 vaccine and how many doses you need.

Off-label use

“Off-label use” means the vaccine is used differently than the way it was originally approved. Vaccine experts support the following off-label uses for KP.2 mRNA COVID-19 vaccines and have no safety concerns:

- You are age 5 years or older and get a dose of Moderna KP.2 vaccine sooner than 6 months after your last dose.
- You are age 5 years or older and get more than 1 dose of a KP.2 vaccine.
- You are age 6 months to 4 years and you get more than 2 doses of a KP.2 vaccine.
- You are age 6 months to 4 years and have already had 2 or more doses of a non-KP.2 vaccine, and you get a dose of a KP.2 vaccine sooner than 6 months after your last dose.

See reverse for more information.

Can I get an mRNA COVID-19 vaccine if I am pregnant or breastfeeding/chestfeeding?

While you are pregnant, you have a higher risk of getting very sick from COVID-19. Getting a COVID-19 vaccine lowers your risk of getting seriously ill from the virus.

You can get an mRNA COVID-19 vaccine if you are pregnant or breastfeeding/chestfeeding. There is more to learn about the Moderna KP.2 and Pfizer-BioNTech KP.2 vaccines while pregnant or breastfeeding/chestfeeding, but the information from the original mRNA vaccines found no concerns. Research shows that mRNA vaccines are the safest type of COVID-19 vaccines to get while pregnant or breastfeeding/chestfeeding.

If you are pregnant or breastfeeding/chestfeeding and have questions about getting a COVID-19 vaccine, talk to your healthcare provider.

Where can I get a COVID-19 vaccine?

Go to ahs.ca/covidvaccine to find out where and when you can get a COVID-19 vaccine. COVID-19 vaccines are free.

Are there side effects from mRNA COVID-19 vaccines?

There can be side effects from mRNA COVID-19 vaccines, but they tend to be mild and go away in a few days. Side effects may include:

- redness, swelling, a hard spot, or feeling sore where you had the needle
- feeling tired or have a headache
- a fever or chills
- body aches or sore joints
- feeling stiff
- pain in your arms or legs
- feeling sick to your stomach (nausea), vomiting (throwing up), loose stool (diarrhea)
- swollen lymph nodes
- swelling or feeling sore in your armpit or groin
- feeling dizzy
- a reduced sense of touch or a feeling of numbness

Children age 5 years and younger may also get an earache, get upset easily, be sleepy, cry, or may not want to eat.

At least 1 out of 100 people who got this vaccine reported 1 or more of these side effects. In some cases, it is unknown if the vaccine caused these side effects.

Current information shows that there is similar risk of side effects after each dose of an mRNA COVID-19 vaccine.

It is important to stay at the clinic for 15 minutes after your vaccine. Some people may have a rare but serious allergic reaction called anaphylaxis. If anaphylaxis happens, you will get medicine to treat the symptoms. It is rare to have a serious side effect. Call Health Link at 811 to report any serious or unusual side effects.

What rare events have been reported after getting an mRNA COVID-19 vaccine?

There have been very rare reports of myocarditis (inflammation of the heart muscle) and pericarditis (inflammation of the lining around the heart) within 7 days of getting an mRNA vaccine. Most reported cases were mild and got better with treatment. The inflammation can cause shortness of breath, chest pain or pressure, or a very fast or abnormal heart rate. Get medical help **right away** if you have any of these symptoms.

Research shows that the risk of these rare events after additional doses is lower than the risk after the second dose for any type of mRNA vaccine. It is expected that the risk will be lower due to the 1-dose schedule for most people. Research is still happening to learn more about the risk of these events after getting updated versions of the mRNA COVID-19 vaccines.

It is not known if having a history of myocarditis or pericarditis puts you at higher risk of having these rare events after a COVID-19 vaccine. Talk to your doctor before you get a dose of COVID-19 vaccine if:

- You had myocarditis or pericarditis within 6 weeks of getting a COVID-19 vaccine.
- You have a history of myocarditis or pericarditis and you have questions about getting a COVID-19 vaccine.

Your risk of getting seriously ill from COVID-19 is much higher than your risk of having a rare event after these vaccines.

How can I manage side effects?

- To help with soreness and swelling, put a cool, wet cloth over the area where you had the needle.
- There is medicine to help with a fever or pain. Check with your doctor or pharmacist if you are not sure what medicine or dose to take. Follow the directions on the package.
- Some people with health problems, such as a weak immune system, must call their doctor if they get a fever. If you have been told to do this, call your doctor even if you think the fever is from the vaccine.

What if I had or am getting another type of vaccine?

You can get most other vaccines at the same time as, any time before, or any time after an mRNA COVID-19 vaccine, including routine vaccines and the influenza vaccine. However, certain vaccines have a waiting period. If you had another vaccine in the last 2 weeks, check with your healthcare provider about when you can get the mRNA COVID-19 vaccine.

Who should not get an mRNA COVID-19 vaccine?

You may not be able to get an mRNA COVID-19 vaccine if:

- You have an allergy to any part of the vaccine.
- You had a severe (serious) or unusual side effect after this vaccine or one like it.
- You are under age 6 months.

If you have allergies or have had a side effect from this vaccine, check with your doctor or a public health nurse before you get the vaccine.

Although you can get the vaccine if you have a mild illness such as a cold or fever, you should stay home until you are feeling better to prevent spreading your illness to others.

Check with your healthcare provider about when you can get an mRNA COVID-19 vaccine if:

- You had COVID-19 in the past.
- You are under age 18 years and have a history of multisystem inflammatory syndrome (MIS-C).

Be sure to talk to your doctor **before** you get an mRNA COVID-19 vaccine if:

- You have a weak immune system (because of a medicine you take or a health problem).
- You have had a stem cell or organ transplant.
- You are getting CAR T-cell therapy.
- You have a history of myocarditis or pericarditis within 6 weeks of getting a dose of COVID-19 vaccine.

For more information about immunization



Call Health Link at 811



Scan the QR code or go to ahs.ca/immunize